

Boundary scattering, symmetric spaces and the principal chiral model on the half-line

N.J. MacKay¹, B. J. Short²

*Department of Mathematics,
University of York,
York YO10 5DD, U.K.*

Abstract

We investigate integrable boundary conditions (BCs) for the principal chiral model on the half-line, and rational solutions of the boundary Yang-Baxter equation (BYBE). In each case we find a connection with (type I, Riemannian, globally) symmetric spaces G/H : there is a class of integrable BCs in which the boundary field is restricted to lie in a coset of H ; these BCs are parametrized by $G/H \times G/H$; there are rational solutions of the BYBE in the defining representations of all classical G parametrized by G/H ; and using these we propose boundary S -matrices for the principal chiral model, parametrized by $G/H \times G/H$, which correspond to our boundary conditions.

¹nm15@york.ac.uk

²bjs108@york.ac.uk

1 Introduction

The bulk principal chiral model (PCM) – that is, the $1 + 1$ -dimensional, $G \times G$ -invariant nonlinear sigma model with target space a compact Lie group G – is known to have a massive spectrum of particles in multiplets which are (sometimes reducible) representations of $G \times G$. These are irreducible representations, however, of the Yangian algebra of non-local conserved charges, and the multiplets are also distinguished by a set of local, commuting conserved charges with spins equal to the exponents of G modulo its Coxeter number. The corresponding bulk scattering (‘ S -’) matrices are constructed from G -invariant (rational) solutions of the Yang-Baxter equation (YBE), whose poles determine the couplings between the multiplets.

In this paper we investigate the model on the half-line – that is, with a boundary. Any proposed boundary S -matrices must satisfy the boundary Yang-Baxter equation (BYBE), for which only a limited range of solutions is known ([7, 8, 9, 11, 12, 13] is a selection). For constant solutions of the BYBE (*i.e.* without dependence on a spectral parameter or rapidity) there is a well-established connection with (quantum) symmetric spaces [37].

We shall find a class of BYBE solutions corresponding to, and parametrized by, the symmetric spaces G/H . These solutions utilize for the bulk S -matrix the rational³, G -invariant solution of the (usual, bulk) YBE in the defining (N -dimensional vector) representation of a classical G , and are themselves rational and N -dimensional, describing the scattering of the bulk vector particle off the boundary ground state. We make the ansatz that they are constant or linear in rapidity, and thus have at most two channels. The underlying algebraic structures are the twisted Yangians [38], though the relationship remains to be explored.

However, we begin by investigating how our solutions might arise as boundary S -matrices, by discussing the principal chiral field on the half-line and boundary conditions which preserve its integrability (see also [31], and [10] for a more general discussion of boundary integrability). We shall find two classes of BCs which are associated with the G/H . In the first, ‘chiral’ class, the field takes its values at the boundary in a coset of H , and the space of such cosets is (up to a discrete ambiguity) $\frac{G}{H} \times \frac{G}{H}$. Correspondingly, we use our BYBE solutions to construct boundary S -matrices, parametrized by $\frac{G}{H} \times \frac{G}{H}$, which preserve the same remnant of the $G \times G$ symmetry as the integrable boundary conditions. In the second, ‘non-chiral’ class, for which we do not generally have corresponding boundary S -matrices, the boundary field lies in a translate of the Cartan immersion of G/H in G . To

³before the inclusion of scalar prefactors

summarize: a connection between boundary integrability and symmetric spaces emerges naturally in two very different ways: by seeking classically integrable boundary conditions, and by solving the BYBE.

The plan of the paper is as follows. In section two, building naturally on the results of [3, 4] for the bulk PCM, we discuss boundary conditions which lead naturally to conservation of local charges. As mentioned, there are two classes of BC, which we call ‘chiral’ and ‘non-chiral’. In section three we find minimal boundary S -matrices, by making ansätze for the BYBE solutions and applying the conditions of crossing-unitarity, hermitian analyticity and R -matrix unitarity, and explain how these are related to symmetric spaces. This section is necessarily rather long and involved, and many of the details appear in appendices. From these, in section four, we construct boundary S -matrices for the PCM, and find that these correspond naturally to the chiral BCs. The key statements of our results for the boundary S -matrices can be found in section 3.4 (for the minimal case, without physical strip poles) and section 4.2 (for the full PCM S -matrices). This paper supersedes the preliminary work of [6].

2 The principal chiral model on the half-line

2.1 The principal chiral model on the full line

We first describe the model on the full line, without boundary. This subsection is largely drawn from [4], and full details may be found there.

The principal chiral model may be defined by the lagrangian

$$\mathcal{L} = \frac{1}{2} \text{Tr} (\partial_\mu g^{-1} \partial^\mu g) , \quad (2.1)$$

where the field $g(x^\mu)$ takes values in a compact Lie group G . (We could also include an overall, coupling constant, but this may be absorbed into $\hbar$, and will not be important for our purposes.) It has a global $G_L \times G_R$ symmetry $g \mapsto g_L g g_R^{-1}$ associated with conserved currents

$$j(x, t)_\mu^L = \partial_\mu g g^{-1}, \quad j(x, t)_\mu^R = -g^{-1} \partial_\mu g \quad (2.2)$$

which take values in the Lie algebra $\mathfrak{g}$ of G : that is, $j = j^a t^a$ (for j^L or j^R : henceforth we drop this superscript) where t^a are the generators of $\mathfrak{g}$, and (with G compact) $\text{Tr}(t^a t^b) = -\delta^{ab}$. The equations of motion are

$$\partial^\mu j_\mu(x, t) = 0, \quad \partial_\mu j_\nu - \partial_\nu j_\mu - [j_\mu, j_\nu] = 0, \quad (2.3)$$

which may be combined as

$$\partial_- j_+ = -\partial_+ j_- = -\frac{1}{2}[j_+, j_-] \quad (2.4)$$

in light-cone coordinates $x^\pm = \frac{1}{2}(t \pm x)$ (and thus $\partial_\pm = \partial_0 \pm \partial_1$).

In addition to the usual spatial parity $P : x \mapsto -x$, the PCM lagrangian has further involutive discrete symmetries. The first, which we call G -parity and which exchanges $L \leftrightarrow R$, is

$$\pi : g \mapsto g^{-1} \Rightarrow j^L \leftrightarrow j^R. \quad (2.5)$$

(In the usual QCD effective model, ‘parity’ is the combination $P\pi$.) Then there is $g \mapsto \alpha(g)$ where α is any involutive automorphism, though only for outer automorphisms may this have a non-trivial effect on the invariant tensors and local charges which we shall consider shortly.

The canonical Poisson brackets for the model are

$$\begin{aligned} \{j_0^a(x), j_0^b(y)\} &= f^{abc} j_0^c(x) \delta(x-y) \\ \{j_0^a(x), j_1^b(y)\} &= f^{abc} j_1^c(x) \delta(x-y) + \delta^{ab} \delta'(x-y) \\ \{j_1^a(x), j_1^b(y)\} &= 0 \end{aligned} \quad (2.6)$$

at equal time. These expressions hold for *either* of the currents j^L or j^R separately, while the algebra of j^L with j^R (which we shall not need here) involves only $\delta'(x-y)$ terms in the brackets of space- with time-components.

This model has two distinct sets of conserved charges, and the two sets commute. The first is the extension of the $G_L \times G_R$ charges to the larger algebra of non-local, Yangian charges [14, 15] $Y(\mathbf{g}_L) \times Y(\mathbf{g}_R)$; we shall not discuss these here. There is also⁴ an infinite set of local, commuting charges with spins s equal to the exponents of $\mathbf{g}$ modulo its Coxeter number,

$$q_{\pm s} = \int_{-\infty}^{\infty} k_{a_1 a_2 \dots a_n} j_{\pm}^{a_1}(x) j_{\pm}^{a_2}(x) \dots j_{\pm}^{a_n}(x) dx \quad (2.7)$$

(where $n = s + 1$); here, unlike for the Yangian, j^L and j^R give the same charges (up to a change of sign), of which there is therefore only one set. The primitive invariant tensors k have to be very carefully chosen to ensure the charges commute – for the full story see [4]. Such a set appears to be precisely what is needed for quantum integrability, where it leads to the beautiful structure of masses and interactions described in [16].

⁴In this paper we restrict to the classical $\mathbf{g}$, although they also exist for exceptional $\mathbf{g}$ [5].

2.2 Boundary conditions for the model on the half-line

Varying the bulk action on the half-line $-\infty < x \leq 0$ imposes the additional boundary equation

$$\text{Tr}(g^{-1}\partial_1 g \cdot g^{-1}\delta g) = 0 \quad \text{at } x = 0,$$

where the variation is over all δg such that $g^{-1}\delta g \in \mathfrak{g}$. Clearly the Neumann condition $\partial_1 g|_0 = 0$ solves this, as does the Dirichlet condition $\delta g|_0 = 0$, or $\partial_0 g|_0 = 0$. But we can also impose mixed conditions, in any way such that

$$(g^{-1}\partial_1 g)^a (g^{-1}\partial_0 g)^a = 0$$

(with the usual summation convention).

We begin by considering some simple mixed boundary conditions written in terms of the currents j . A little later we shall generalize these, and write them in terms of the fields g . We take as a BC on the currents

$$j_+^a(0) = R^{ab} j_-^b(0), \tag{2.8}$$

with each j chosen independently to be either L or R; we refer to the four possibilities as LL, RR, LR and RL. The boundary equation of motion then requires that R^{ab} be an orthogonal matrix. We would also like consistency with the Poisson brackets: if we extend the currents' domains to $x > 0$ by requiring $j_+^a(x) = R^{ab} j_-^b(-x)$, then this further requires that R be symmetric, and give an (involutive; $\alpha^2 = 1$) automorphism α of $\mathfrak{g}$ via

$$\alpha : t^a \mapsto R^{ab} t^b.$$

Together these imply that R is diagonalizable with eigenvalues ± 1 , so that we may write

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{k},$$

where $\mathfrak{h}$ and $\mathfrak{k}$ are the $+1$ and -1 eigenspaces respectively. The $\mathfrak{h}$ indices then correspond to Neumann directions $j_+^a = j_-^a \Rightarrow j_1^a = 0$, the $\mathfrak{k}$ indices to Dirichlet directions $j_+^a = -j_-^a \Rightarrow j_0^a = 0$ (all at $x = 0$). Further, R 's being an automorphism implies that

$$[\mathfrak{h}, \mathfrak{h}] \subset \mathfrak{h}, \quad [\mathfrak{h}, \mathfrak{k}] \subset \mathfrak{k}, \quad [\mathfrak{k}, \mathfrak{k}] \subset \mathfrak{h},$$

precisely the properties required of a symmetric space G/H [17], where H is the subgroup generated by $\mathfrak{h}$ and invariant under the involution α .

For integrability we require a great deal of R . As we have said, there are two infinite sets of charges. The Yangian charges appear no longer to be conserved on the half-line, even

with pure Neumann BCs [19] (naively, at least, it seems that there are remnants only, as we shall see). However, we believe they are not essential for integrability, because precisely half of the local charges remain conserved, with either $q_s + q_{-s}$ or $q_s - q_{-s}$ surviving. Our conjecture is that these are enough to guarantee the properties of quantum integrability, such as factorizability of the S -matrix.

The first charge is energy, and its conservation on the half-line is just the equation of motion, requiring R to be orthogonal. For the higher charges we take

$$\begin{aligned} q_{|s|} &= q_s \pm q_{-s} \\ &= \int_{-\infty}^0 k_{a_1 a_2 \dots a_n} \{ j_+^{a_1} \dots j_+^{a_n} \pm j_-^{a_1} \dots j_-^{a_n} \} dx. \end{aligned}$$

Then

$$\begin{aligned} \frac{d}{dt} q_{|s|} &= \int_{-\infty}^0 k_{a_1 a_2 \dots a_n} \{ (\partial_- + \partial_1) (j_+^{a_1} \dots j_+^{a_n}) \pm (\partial_+ - \partial_1) (j_-^{a_1} \dots j_-^{a_n}) \} dx \\ &= k_{a_1 a_2 \dots a_n} \{ j_+^{a_1} \dots j_+^{a_n} \mp j_-^{a_1} \dots j_-^{a_n} \} |_{x=0} \\ &= (k_{b_1 b_2 \dots b_n} R^{a_1 b_1} \dots R^{a_n b_n} \mp k_{a_1 \dots a_n}) j_-^{a_1} \dots j_-^{a_n} |_{x=0}. \end{aligned}$$

That this is zero for one choice of sign follows from the result [20] that, for every R and k ,

$$k_{b_1 b_2 \dots b_n} R^{a_1 b_1} \dots R^{a_n b_n} = \epsilon k_{a_1 \dots a_n}, \quad (2.9)$$

where $\epsilon = \pm 1$. (This is obvious, with $\epsilon = 1$, when α is an inner automorphism, but not at all obvious for outer automorphisms.) So, if we now regard α as acting on the currents, $\alpha(j_\pm^a) = R^{ab} j_\pm^b$, we see that

$$q_{|s|} = \int_{-\infty}^0 k_{a_1 a_2 \dots a_n} \{ j_+^{a_1} \dots j_+^{a_n} + \alpha(j_-^{a_1}) \dots \alpha(j_-^{a_n}) \} dx \quad (2.10)$$

is the charge which remains conserved in the presence of the boundary. For LL and RR conditions its density is the combination which is invariant under the combined action $P\alpha$ of spatial parity P (which exchanges $j_+ \leftrightarrow j_-$) and α , while for LR and RL it is the combination invariant under these together with G -parity, $P\alpha\pi$.

Further, these charges still commute. In the Poisson bracket of the charges constructed from tensors $k^{(1)}$ and $k^{(2)}$, a total derivative term which vanished in the bulk now gives an additional contribution proportional to (all at $x = 0$)

$$\begin{aligned} & k_{ca_1 \dots a_s}^{(1)} k_{cb_1 \dots b_r}^{(2)} (j_+^{a_1} \dots j_+^{a_s} j_+^{b_1} \dots j_+^{b_r} - \epsilon^{(1)} \epsilon^{(2)} j_-^{a_1} \dots j_-^{a_s} j_-^{b_1} \dots j_-^{b_r}) \\ &= k_{cd_1 \dots d_s}^{(1)} k_{ce_1 \dots e_r}^{(2)} (R^{d_1 a_1} \dots R^{d_s a_s} R^{e_1 b_1} \dots R^{e_r b_r} - \epsilon^{(1)} \epsilon^{(2)} \delta^{d_1 a_1} \dots \delta^{d_s a_s} \delta^{e_1 b_1} \dots \delta^{e_r b_r}) j_-^{a_1} \dots j_-^{a_s} j_-^{b_1} \dots j_-^{b_r} \\ &= k_{d_0 d_1 \dots d_s}^{(1)} k_{e_0 e_1 \dots e_r}^{(2)} (R^{d_0 a_0} R^{d_1 a_1} \dots R^{e_0 b_0} R^{e_1 b_1} \dots - \epsilon^{(1)} \epsilon^{(2)} \delta^{d_0 a_0} \delta^{d_1 a_1} \dots \delta^{e_0 b_0} \delta^{e_1 b_1} \dots) \delta^{a_0 b_0} j_-^{a_1} \dots j_-^{b_r} \\ &= 0 \end{aligned}$$

by property (2.9). Finally, it is precisely the $q_{|s|}$ of (2.10) above that still commute with the $G \times G$ -generating charges $Q = \int_{-\infty}^0 j_0^{L/R} dx$.

2.2.1 General chiral BCs

As we commented when first introducing the BC (2.8), in $j_+ = \alpha(j_-)$ we may take each j as either L or R. The LL and RR conditions are then related: $j_+^L = \alpha(j_-^L) \Rightarrow gj_+^R g^{-1} = \alpha(gj_-^R g^{-1})$ (all at $x = 0$). In fact the most general such (we shall call it ‘chiral’) BC is to take

$$g(0) \in k_L H k_R^{-1}. \quad (2.11)$$

This is the Dirichlet part of the BC; when we impose the boundary equation-of-motion we supplement it with Neumann conditions within this boundary target space (which we shall henceforth refer to as the D-submanifold) so that the current conditions become

$$\begin{aligned} k_L^{-1} j_+^L k_L &= \alpha(k_L^{-1} j_-^L k_L) \\ k_R^{-1} j_+^R k_R &= \alpha(k_R^{-1} j_-^R k_R) \quad (\text{at } x = 0). \end{aligned}$$

The constant group elements k_L and k_R parametrize left- and right-cosets of H in G and may be taken to lie in the Cartan immersion of G/H in G , so that the possible BCs are parametrized by $G/H \times G/H$. (In fact this is true only at the level of the Lie algebras: there is a further discrete ambiguity in the choice of k_L, k_R . For details of this, and of the Cartan immersion, we refer the reader to appendices 6.1, 5.1.) Our earlier results about conservation and commutation of charges and consistency with the Poisson brackets still apply (generalized here by twisting the currents with an inner automorphism, which does not change the definition of the conserved charge $q_{|s|}$).

Note that when $k_L = k_R = e$ (where e is the identity element in G), we have $g(0) \in H$, the continuous Dirichlet boundary parameters which determine the D-submanifold are all trivial, and the residual symmetry is $H \times H$. For any k_L, k_R the case $H = G$ corresponds to the pure Neumann condition, while trivial H , the pure Dirichlet condition, is inadmissible for any non-abelian G .

We should point out at this stage that we have not succeeded in finding a boundary Lagrangian for any of our mixed BCs. That is, we have no Lagrangian of which the free variation leads to our conditions. The Dirichlet conditions have to be imposed as ‘clamped’ BCs, restricting the boundary variation of g .

Let us now examine how much of the $G \times G$ symmetry survives. We can see that the BC (2.11) is invariant under $k_L H k_L^{-1} \times k_R H k_R^{-1}$, and we can check that it is precisely the

charges generating this subgroup of $G \times G$ which are conserved on the half-line. For the global G -generating charges $Q^a = \int_{-\infty}^0 j_0^a(x) dx$ (where subscripts either all L or all R are to be understood), consider

$$\begin{aligned} \frac{d}{dt} (k^{-1} Q k) &= k^{-1} \left(\int_{-\infty}^0 \partial_0 j_0 dx \right) k \\ &= k^{-1} j_1(0) k \\ &= \frac{1}{2} [k^{-1} j_+ k - k^{-1} j_- k]_{x=0} \\ &= \frac{1}{2} [\alpha(k^{-1} j_- k) - k^{-1} j_- k]_{x=0} , \end{aligned}$$

which is zero on $\mathfrak{h}$ (only).

2.2.2 General non-chiral BCs

If we explore similarly the LR and RL conditions, we find that the condition

$$g(0) \in g_L \frac{G}{H} g_R^{-1} \quad (2.12)$$

(where $G/H = \{\alpha(g)g^{-1} | g \in G\}$ is the Cartan immersion of G/H in G) leads to $g_L^{-1} j_0^L g_L = \alpha(g_R^{-1} j_0^R g_R)$ (with g_L, g_R again constant elements of G). If we then apply the boundary equation-of-motion, the condition on the currents becomes

$$g_L^{-1} j_{\pm}^L g_L = \alpha(g_R^{-1} j_{\mp}^R g_R) \quad \text{at } x = 0.$$

Unlike our chiral BCs (which are parametrized by $G/H \times G/H$), these BCs are parametrized by a single G (again quotiented by a discrete subgroup; see appendix 6.2). In the specialization $g_L = g_R$, however, the boundary is parametrized by G/H .

Note the inversion of the role of the dimension of H in determining the dimension of the D-submanifold, compared to the chiral case: there (and setting $k_L = k_R = e$) we had $g(0) \in H$, whereas here (with $g_L = g_R = e$) we have $g(0) \in G/H$. The two extreme non-chiral cases give us nothing new: with trivial H we revert to the free, pure Neumann condition, while at the other extreme of $H = G$ we have the pure Dirichlet condition.

As with the chiral case, we can check conservation of the generators of the remnant of the $G \times G$ symmetry. This time, because the Cartan immersion of G/H is invariant under H_{diag} (the diagonal subgroup $g \mapsto hgh^{-1}$), the surviving global symmetry is H_{diag} conjugated (in $G \times G$) by (g_L, g_R) , and we may check that

$$\frac{d}{dt} (g_L^{-1} Q_L g_L + g_R^{-1} Q_R g_R) = \frac{1}{2} [g_L^{-1} (j_+^L - j_-^L) g_L + g_R^{-1} (j_+^R - j_-^R) g_R]_{x=0}$$

$$= \frac{1}{2} [g_R^{-1}(j_+^R - j_-^R)g_R - \alpha(g_R^{-1}(j_+^R - j_-^R)g_R)]_{x=0}$$

which is zero precisely on $\mathbf{h} \times \mathbf{h}$.

At this stage it is worth comparing our results with those obtained in the Wess-Zumino-Witten model – that is, with D-branes on group manifolds. There, initial suggestions of a connection with symmetric spaces [21] were supplanted by an understanding that the D-submanifold is actually a ‘twisted’ or ‘twined’ conjugacy class [22, 23], $C_\alpha(g_0) = \{\alpha(g)g_0g^{-1} | g \in G\}$. This situation arises because in the WZW model there is only one pair of currents, j_+^L and j_-^R , and so only one, LR, boundary condition. In our case we have two, LR and RL, conditions, and their interplay further requires that $\alpha(g_0) = g_0^{-1}$. But the space $\mathcal{M}$ of such g_0 is, for the non-Grassmannian cases, precisely the Cartan immersion G/H (which is connected to the identity, so that we can find $\sqrt{g_0}$), and

$$C_\alpha(g_0) = \{\alpha(g)g_0g^{-1} | g \in G\} = \{\alpha(gg_0^{-\frac{1}{2}})(gg_0^{-\frac{1}{2}})^{-1} | g \in G\} = \frac{G}{H}. \quad (2.13)$$

(For the Grassmannian cases, $\mathcal{M}$ is a union of disconnected components, the identity-connected component being G/H – see appendix 5. However, each of the other components is actually a translate of the immersion of G/H' for a different H' [40], so we obtain no new BCs in this way.)

The analogous BC in our case is with $g_L = g_R = e$, and the residual symmetry is $H_{\text{diag.}}$, necessarily preserved by any BC utilizing a twisted conjugacy class, as in the WZW model. Note that the $\alpha = 1, H = G$ case is purely Dirichlet in our case, whereas in the WZW model g_0 is unconstrained and there is still freedom at the boundary. We expect that the non-chiral BCs should remain integrable when a Wess-Zumino term (of arbitrary size) is added, and we plan to explore this in future work.

Finally we note the relationship of our work to that on the Gross-Neveu model⁵ [25]. This model has a single global $G = O(N)$ invariance, broken by the BC to an $H = O(M)$ subgroup. Their boundary S -matrix is then diagonal.

2.3 Remarks on quantization

As with the bulk model, we shall assume that our results carry through into the quantum theory: that classically conserved charges remain conserved in the quantum theory; that charges classically in involution do not develop $\mathcal{O}(\hbar^2)$ anomalies in their commutators; and that our BCs therefore lead to quantum conservation of the charges which generate

⁵not the generalized chiral GN model, which remains to be investigated

the residues of the $G \times G$ symmetry. All of this leads to the expectation that boundary scattering factorizes, so that solutions of the BYBE provide boundary S -matrices.

The only technique which can give evidence for the continued conservation of the local charges after quantization is Goldschmidt-Witten anomaly counting [26]. This was carried out for the bulk case in [3], where for each classical G at least one non-trivial charge was found to be necessarily conserved in the quantum theory. This was extended to boundary models in [24], and used to prove quantum conservation of the spin-3 charge for $G = SO(N)$ in [6] for one of our BCs. It is simple to check that for all our BCs and for all classical G , each charge which necessarily survives quantization in the bulk model also survives in the presence of the boundary. We do not give details.

Finally, the form of our admissible D-submanifolds is not so surprising when we remember that bulk sigma models on symmetric spaces have particularly nice behaviour after quantization, in that the symmetric spaces preserve their shape under renormalization [41]. We would certainly expect our D-submanifolds to behave similarly nicely.

3 The minimal boundary S -matrices

In this section we construct boundary S -matrices which are minimal — that is, which have no poles on the physical strip. We follow the method used in the bulk case [29], where minimal S -matrices were found by solving the Yang-Baxter equation and applying unitarity, analyticity and crossing symmetry, and the desired pole structure then implemented using the CDD ambiguity. In the boundary case we seek minimal boundary S -matrices by making ansätze to solve the boundary Yang-Baxter equation (BYBE) and applying unitarity, analyticity and the combined crossing-unitarity relation [27]. These minimal solutions will be used to construct PCM boundary S -matrices with the appropriate pole structure in section four.

We shall make the ansatz that the boundary S -matrix (the ‘ K -matrix’) in the defining, N -dimensional, vector representation of a classical group G is in one of the two forms

$$K_1(\theta) = \rho(\theta)E \quad \text{and} \quad K_2(\theta) = \frac{\tau(\theta)}{(1 - c\theta)}(I + c\theta E). \quad (3.1)$$

Here c and E are constants, the latter an $N \times N$ matrix; we shall explain the θ -dependent terms below. The crucial point at this stage is the equivalent physical statement that K has at most two ‘channels’: since its matrix structure is at most linear in rapidity θ , it will decompose into one (for K_1) or two (for K_2) projectors. This will prove sufficient to yield a set of solutions related to the symmetric spaces in the following


Correspondence: for a given G -invariant bulk factorized S -matrix (*i.e.* a solution of the YBE) in the defining representation, the K -matrices of the form (3.1) fall into a set of families in 1–1 correspondence with the set of symmetric spaces G/H , and each family is parametrized by a space of admissible E which is isomorphic to (possibly a finite multiple of) the corresponding G/H .

Such solutions, we believe, correspond to scattering off the boundary ground state. We would obtain solutions with many more channels by considering scattering of bulk particles in higher tensor representations or off higher boundary bound states, or both. These can be obtained by fusion from our solutions⁶, and the results of this paper thus lay the foundation for future work in this direction.

The calculations of this section will necessarily, because case-by-case and exhaustive, be rather involved. Our strategy is to lead the reader through the implications of the BYBE, unitarity, hermitian analyticity and crossing-unitarity, initially culminating in a precise statement of our solutions in section 3.4. Details of the calculations are relegated to appendices 7 and 8. Then in section 3.5 we explain how our solutions are parametrized by the symmetric spaces; details appear in appendix 5.

3.1 Calculating the boundary S -matrices

Throughout the rest of this paper we shall use the following (somewhat unconventional) notation for the bulk and boundary S -matrices:



where θ is the rapidity difference between the two in-coming particles which scatter in the first diagram and ϕ is the rapidity of the in-coming particle reflecting in the second.

We consider the BYBE for two particles in the vector representation

$$S_{ij}^{kl}(\theta - \phi)(I_{jm} \otimes K^{ln}(\theta))S_{mo}^{np}(\theta + \phi)(I_{oq} \otimes K^{pr}(\phi)) = \\ (I_{ij} \otimes K^{kl}(\phi))S_{jm}^{ln}(\theta + \phi)(I_{mo} \otimes K^{np}(\theta))S_{oq}^{pr}(\theta - \phi) \quad (3.2)$$

⁶with the exception of the $SO(N)$ spinorial multiplets

(θ and ϕ are now the rapidities of the two particles.) We attempt to find solutions of the form (equivalent to (3.1))

$$K_1(\theta) = \rho(\theta)E \quad \text{and} \quad K_2(\theta) = \tau(\theta) \left(P^- - \left[\frac{h}{ci\pi} \right] P^+ \right) \quad (3.3)$$

where $\rho(\theta)$ and $\tau(\theta)$ are scalar prefactors, h is the dual Coxeter number of the group, c is a parameter and

$$P^\pm = \frac{1}{2} (I \pm E) \quad [x] = \frac{\theta + \frac{i\pi x}{h}}{\theta - \frac{i\pi x}{h}},$$

where I is the identity matrix and E is a general square matrix of the same size⁷. We find that constraints are imposed on the scalar prefactors and the matrix E by the BYBE and unitarity, analyticity and crossing-unitarity. These constraints are in general dependent on the choice of classical group, $SU(N)$, $SO(N)$ or $Sp(N)$, but in all cases involving $K_2(\theta)$ we find that $E^2 = 1$, so that $P^\pm$ are projectors, as the notation suggests.

We note that the constraints imposed on the scalar prefactors will allow us to find them only up to the usual CDD ambiguity. This freedom is then further restricted by taking the K -matrices to be minimal – that is, taking $\rho(\theta)$ and $\tau(\theta)$ to have no poles on the physical strip. We shall also require $\tau(\theta)$ to have a zero at $\theta = \frac{1}{c}$ so that $K_2(\theta)$ is finite at this point, since we shall find that $\frac{1}{c}$ can lie in the physical strip.

For the case of $SU(N)$ the vector representation is not self-conjugate. This allows us to consider the situation where a particle scattering off the boundary returns as an anti-particle (for an analogous situation see [11]). In this case the K -matrix, $K^{i\bar{j}}(\theta)$, must satisfy a version of the BYBE which for convenience we shall refer to as the ‘conjugated’ BYBE,

$$\begin{aligned} S_{ij}^{kl}(\theta - \phi)(I_{jm} \otimes K^{l\bar{n}}(\theta))S_{m\bar{o}}^{\bar{n}p}(\theta + \phi)(I_{\bar{o}\bar{q}} \otimes K^{p\bar{r}}(\phi)) = \\ (I_{ij} \otimes K^{k\bar{l}}(\phi))S_{j\bar{m}}^{\bar{l}n}(\theta + \phi)(I_{\bar{m}\bar{o}} \otimes K^{n\bar{p}}(\theta))S_{\bar{o}\bar{q}}^{\bar{p}\bar{r}}(\theta - \phi). \end{aligned} \quad (3.4)$$

We shall find that only $K_1(\theta)$ gives a solution of the conjugated BYBE. Analyticity and unitarity become more subtle in this case. Henceforth we shall refer to this case as the ‘conjugating’ case for the $SU(N)$ model.

Even in the ‘non-conjugating’ $SU(N)$ model, where the representation is preserved under scattering by the boundary, the fact that the vector representation is not self-conjugate leads to subtleties in the calculations. In addition to the standard BYBE of (3.2) there is also the possibility that one or both of the in-coming particles in the BYBE scattering

⁷The apparently circuitous involvement of $i\pi/h$ ensures that x is an integer, as is usual in the literature.

process is the conjugate vector particle. Thus we need to introduce a minimal K -matrix, $\bar{K}(\theta)$, describing this scattering. We take $\bar{K}(\theta)$ to have the same one or two channel form as $K(\theta)$, that is

$$\bar{K}_1(\theta) = \bar{\rho}(\theta)F \quad \text{or} \quad \bar{K}_2(\theta) = \frac{\bar{\tau}(\theta)}{(1 - d\theta)}(I + d\theta F), \quad (3.5)$$

where we have a new set of scalar prefactors and constants, $\bar{\rho}(\theta)$, $\bar{\tau}(\theta)$, d and F .

At this point we introduce the diagrammatic algebra used in the calculations we perform. We represent the matrices I , J , E and F , where J is the symplectic form matrix, in the following way:

$$I: \text{---} \quad J: \rightarrow \quad E: \text{---} \circ \quad F: \bullet \text{---} \quad \left(J^t: \leftarrow \right)$$

(Thus from the properties of J we have $\rightarrow = -(\leftarrow)$ and $\rightarrow \leftarrow = \text{---} = \leftarrow \rightarrow$.) Matrix multiplication is achieved by concatenation of the diagrams, for example

$$JIJ^tE: \rightarrow \leftarrow \circ = \rightarrow \leftarrow \circ = \circ$$

Using this diagrammatic algebra we can rewrite the K -matrices of (3.1) as

$$K_1(\theta) = \rho(\theta) \text{---} \circ \quad K_2(\theta) = \frac{\tau(\theta)}{(1 - c\theta)} \left(\text{---} + c\theta \text{---} \circ \right) \quad (3.6)$$

This diagrammatic description is used in the calculations presented in the appendices, and makes them much clearer.

We substitute either of these K -matrices into the BYBE, taking the minimal S -matrix to be that of the bulk PCM for a particular G , derived in [29] up to a few minor inconsistencies which we have corrected. This yields constraints (dependent on G and on whether we take K_1 or K_2) that E (and F) must satisfy in order for K_i to be a solution of the BYBE. In fact the constraints from crossing-unitarity on the K_i are identical to those imposed by the BYBE, therefore we shall delay presentation of the constraints until later. (The reader is referred to appendix 7.1 for details of the BYBE calculations.) Instead we go on to consider unitarity, analyticity and crossing-unitarity.

3.2 Unitarity and Analyticity

3.2.1 The non-conjugating cases

For the non-conjugating cases the K -matrices are required to satisfy the conditions of unitarity [27] and hermitian analyticity [28]

$$K(\theta)K(-\theta) = I \quad K(\theta) = (K(-\theta^*))^\dagger. \quad (3.7)$$

Substituting K_1 we obtain

$$\rho(\theta)\rho(-\theta)\text{---}\text{---} = \text{---} \quad \rho(\theta)\text{---}\text{---} = \rho(-\theta^*)^*(\text{---}\text{---})^\dagger,$$

These matrix equations are equivalent to

$$\begin{aligned} \rho(\theta)\rho(-\theta) &= \frac{1}{\alpha} & \rho(\theta) &= \beta\rho(-\theta^*)^* \\ \text{---}\text{---} &= \alpha\text{---} & (\text{---}\text{---})^\dagger &= \beta\text{---}\text{---} \end{aligned}$$

where α and β are constants with $\beta \in U(1)$. Recalling $K_1(\theta)$, we see that we have the freedom in our definition of $\rho(\theta)$ and $\text{---}\text{---}$ to set $\beta = 1$ and ensure $\alpha \in U(1)$, so that the constraints imposed on the matrix $\text{---}\text{---}$ by unitarity and analyticity become

$$(\text{---}\text{---})^\dagger = \text{---}\text{---} \quad \text{and} \quad \text{---}\text{---} = \alpha\text{---} \quad \text{where } \alpha \in U(1). \quad (3.8)$$

Similarly for K_2 we obtain

$$\begin{aligned} \frac{\tau(\theta)\tau(-\theta)}{(1-c^2\theta^2)} \left(\text{---} - c^2\theta^2\text{---}\text{---} \right) &= \text{---} \\ \frac{\tau(\theta)}{(1-c\theta)} \left(\text{---} + c\theta\text{---}\text{---} \right) &= \frac{\tau(-\theta^*)^*}{(1+c^*\theta)} \left(\text{---} - c^*\theta(\text{---}\text{---})^\dagger \right) \end{aligned}$$

These are equivalent to

$$\begin{aligned} \tau(\theta)\tau(-\theta) &= \frac{(1-c^2\theta^2)}{(1-\gamma c^2\theta^2)} & \text{---}\text{---} &= \gamma\text{---} \\ \frac{\tau(\theta)}{(1-c\theta)} &= \frac{\tau(-\theta^*)^*}{(1+c^*\theta)} & c\text{---}\text{---} &= -c^*(\text{---}\text{---})^\dagger \end{aligned}$$

If we consider the expression for $K_2(\theta)$, we see that we have the freedom to set c to be purely imaginary and choose $\gamma \in U(1)$. Then the constraints on the matrix $\text{---}\text{---}$ become

$$(\text{---}\text{---})^\dagger = \text{---}\text{---} \quad \text{and} \quad \text{---}\text{---} = \gamma\text{---} \quad \text{where } \gamma \in U(1)$$

as was the case for K_1 . In fact we find that the parameters α and γ cannot be freely chosen in $U(1)$; the only choice consistent with the hermiticity of $\text{---}\text{---}$ is that they are equal to 1 (see appendix 8.1). Consequently we find that for both K_1 and K_2 , unitarity and analyticity impose

$$(\text{---}\text{---})^\dagger = \text{---}\text{---} \quad \text{and} \quad \text{---}\text{---} = \text{---}. \quad (3.9)$$

The corresponding constraints imposed on $\rho(\theta)$ and $\tau(\theta)$ are

$$\rho(\theta) = \rho(-\theta^*)^* \quad \rho(\theta)\rho(-\theta) = 1 \quad (3.10)$$

$$\tau(\theta) = \tau(-\theta^*)^* \quad \tau(\theta)\tau(-\theta) = 1. \quad (3.11)$$

(We note that similar conditions are imposed on $\bar{\rho}(\theta)$, $\bar{\tau}(\theta)$ and $\text{---}\bullet\text{---}$.)

3.2.2 $SU(N)$ -conjugating

In the case of the conjugated BYBE (3.4), which we call ‘ $SU(N)$ -conjugating’, unitarity and analyticity can no longer be applied as straightforwardly. The reason is that we are no longer dealing with a K -matrix that is an endomorphism of the vector representation space V , but rather with $K(\theta) : V \rightarrow \bar{V}$. In order to apply our conditions we must introduce $K'(\theta) : \bar{V} \rightarrow V$ and consider the space $V \oplus \bar{V}$ on which the endomorphism

$$\tilde{K}(\theta) = \begin{pmatrix} 0 & K'(\theta) \\ K(\theta) & 0 \end{pmatrix} \quad \text{acts.}$$

We can then apply analyticity and unitarity to $\tilde{K}(\theta)$ which yields

$$K(\theta) = K'(-\theta^*)^\dagger \quad K(\theta)K'(-\theta) = I. \quad (3.12)$$

The conjugated BYBE that $K(\theta)$ and $K'(\theta)$ must satisfy allows only the $K_1(\theta)$ form for both. Thus we have

$$K(\theta) = \rho(\theta) \text{---}\bigcirc \quad \text{and} \quad K'(\theta) = \omega(\theta) \text{---}\bullet$$

where $\rho(\theta)$ and $\omega(\theta)$ are scalar prefactors, whilst $\text{---}\bigcirc = E$ and $\text{---}\bullet \equiv F$ are constant matrices. Substituting these into our conditions yields

$$\begin{aligned} \rho(\theta) &= \alpha \omega(-\theta^*)^* & \rho(\theta) \omega(-\theta) &= \frac{1}{\beta} \\ \alpha \text{---}\bigcirc &= (-\bullet \text{---})^\dagger & \text{---}\bigcirc \text{---}\bullet &= \beta \text{---} \end{aligned} \quad (3.13)$$

There is enough rescaling freedom in splitting $K(\theta)$ and $K'(\theta)$ into scalar prefactors and matrix parts that we can consistently set $\alpha = \beta = 1$ and $\det E = 1 \iff \det F = 1$ (details in appendix 8.2). This leaves us with

$$\begin{aligned} E &= F^\dagger, & EF &= I, & \det F &= \det E = 1, \\ \rho(\theta) &= \omega(-\theta^*)^* & \text{and} & & \rho(\theta) \omega(-\theta) &= 1. \end{aligned} \quad (3.14)$$

3.3 Crossing-Unitarity

3.3.1 The non-conjugating cases

The boundary S -matrices must also satisfy crossing-unitarity [27], which in our notation is

$$K^{ij}(\frac{i\pi}{2} - \theta) = S_{jk}^{i\bar{l}}(2\theta) K^{\bar{l}k}(\frac{i\pi}{2} + \theta). \quad (3.15)$$

Substituting the K_i into the crossing-unitarity equation gives constraints on $\text{--}\circ\text{--}$, c , d and the scalar prefactors. We tabulate these below (details in appendix 7.2). Note that in our diagrammatic algebra $Tr(E)$ is represented by the symbol $\bigcirc$.

Group $K_1(\theta)/K_2(\theta)$	Constraints
$SU(N)$ $K_1(\theta)$	$\bigcirc = 0$, $\rho(i\pi - \theta)\rho(i\pi + \theta) = \frac{(\theta - \frac{i\pi}{h})(\theta + \frac{i\pi}{h})}{\theta^2}$
$SU(N)$ $K_2(\theta)$	$\bigcirc = \frac{ih}{\pi} (\frac{1}{c} + \frac{1}{d})$, $\tau(i\pi - \theta)\tau(i\pi + \theta) = \frac{(\theta - \frac{i\pi}{h})(\theta + \frac{i\pi}{h})(\theta - i\pi + \frac{1}{c})(\theta + i\pi - \frac{1}{c})}{(\theta - i\pi)(\theta + i\pi)(\theta - \frac{1}{d})(\theta + \frac{1}{d})}$
$SO(N)$ $K_1(\theta)$	$\bigcirc = 0$ and either $(\text{--}\circ\text{--})^t = \text{--}\circ\text{--}, \quad \frac{\rho(i\pi/2 - \theta)}{\rho(i\pi/2 + \theta)} = \sigma_o(2\theta)$ or $(\text{--}\circ\text{--})^t = -(\text{--}\circ\text{--}), \quad \frac{\rho(i\pi/2 - \theta)}{\rho(i\pi/2 + \theta)} = -[1]\sigma_o(2\theta)$
$SO(N)$ $K_2(\theta)$	$\bigcirc = \frac{2ih}{c\pi}, \quad (\text{--}\circ\text{--})^t = \text{--}\circ\text{--}, \quad \frac{\tau(i\pi/2 - \theta)}{\tau(i\pi/2 + \theta)} = [\frac{h}{2}] [\frac{h}{ci\pi} - \frac{h}{2}] \sigma_o(2\theta)$ or $(\text{--}\circ\text{--})^t = -(\text{--}\circ\text{--}), \quad \frac{\tau(i\pi/2 - \theta)}{\tau(i\pi/2 + \theta)} = -[1][\frac{h}{2}] [\frac{h}{ci\pi} - \frac{h}{2}] \sigma_o(2\theta)$
$Sp(N)$ $K_1(\theta)$	$\bigcirc = 0$ and either $\rightarrow(\text{--}\circ\text{--})^t\rightarrow = \text{--}\circ\text{--}, \quad \frac{\rho(i\pi/2 - \theta)}{\rho(i\pi/2 + \theta)} = \sigma_p(2\theta)$ or $\rightarrow(\text{--}\circ\text{--})^t\rightarrow = -(\text{--}\circ\text{--}), \quad \frac{\rho(i\pi/2 - \theta)}{\rho(i\pi/2 + \theta)} = -[1]\sigma_p(2\theta)$
$Sp(N)$ $K_2(\theta)$	$\rightarrow(\text{--}\circ\text{--})^t\rightarrow = \text{--}\circ\text{--}, \quad \frac{\tau(i\pi/2 - \theta)}{\tau(i\pi/2 + \theta)} = [\frac{h}{2}] [\frac{h}{ci\pi} - \frac{h}{2}] \sigma_p(2\theta)$ or $\bigcirc = \frac{2ih}{c\pi}, \quad \rightarrow(\text{--}\circ\text{--})^t\rightarrow = -\text{--}\circ\text{--},$ $\frac{\tau(i\pi/2 - \theta)}{\tau(i\pi/2 + \theta)} = -[1][\frac{h}{2}] [\frac{h}{ci\pi} - \frac{h}{2}] \sigma_p(2\theta)$

The functions σ_o and σ_p are scalar prefactors for the bulk S -matrices (see appendix 7).

3.3.2 $SU(N)$ -conjugating

The $SU(N)$ -conjugating K -matrix must satisfy the crossing-unitarity equation

$$K^{i\bar{j}}(\frac{i\pi}{2} - \theta) = S_{jk}^{il}(2\theta) K^{l\bar{k}}(\frac{i\pi}{2} + \theta). \quad (3.16)$$

From this we obtain the constraints (see appendix 7)

$$\frac{\rho(\frac{i\pi}{2} - \theta)}{\rho(\frac{i\pi}{2} + \theta)} = \begin{cases} \sigma_u(2\theta) & (-\circ-)^t = -\circ- \\ -[1]\sigma_u(2\theta) & (-\circ-)^t = -(-\circ-). \end{cases} \quad (3.17)$$

Similarly, the crossing-unitarity equation for $K'(\theta)$ yields identical constraints on $\omega(\theta)$ and $-\bullet-$. Since $-\circ-\bullet- = -\circ-$ the signs in $(-\circ-)^t = \pm(-\circ-)$ and $(-\bullet-)^t = \pm(-\bullet-)$ must coincide. From this we have

$$\frac{\rho(\frac{i\pi}{2} - \theta)}{\rho(\frac{i\pi}{2} + \theta)} = \frac{\omega(\frac{i\pi}{2} - \theta)}{\omega(\frac{i\pi}{2} + \theta)}, \quad (3.18)$$

which, together with the other constraints on these scalar prefactors, is enough to show we should take $\rho(\theta) = \omega(\theta)$.

Now F and $\omega(\theta)$ are completely fixed by E and $\rho(\theta)$, and the boundary S -matrix for $V \oplus \bar{V}$ is

$$\tilde{K}(\theta) = \rho(\theta) \begin{pmatrix} 0 & E^\dagger \\ E & 0 \end{pmatrix} \quad (3.19)$$

subject to the following constraints, along with (3.17):

$$\begin{aligned} E^\dagger E &= I & \text{and} & & \det E &= 1 \\ \rho(\theta) &= \rho(-\theta^*)^* & \text{and} & & \rho(\theta)\rho(-\theta) &= 1. \end{aligned} \quad (3.20)$$

3.4 The boundary S -matrices

We have obtained a series of constraints on $-\circ-$ and on $\rho(\theta)$ and $\tau(\theta)$ which must be satisfied if the proposed $K_1(\theta)$ and $K_2(\theta)$ are to be boundary S -matrices. The constraints on the scalar prefactors enable us to determine them exactly for each G , providing we assume some extra minimality conditions, namely that they should be meromorphic functions of θ with no poles on the physical strip $\text{Im } \theta \in [0, \frac{\pi}{2}]$. Having calculated the scalar prefactors (we do not include details of the calculations, as the reader can simply check the results if required) we obtain the boundary S -matrices below.

We note that in the case of $K_2(\theta)$ there is also the possibility that the pole at $\theta = \frac{1}{c}$ lies on the physical strip and so the scalar prefactor $\tau(\theta)$ may be required to have a zero at this point. In fact we shall find that one of $\pm \frac{1}{c}$ always lies on the physical strip, and the expressions given below for $K_2(\theta)$, for each case, are valid when $\frac{1}{c}$ lies on the physical strip. When $-\frac{1}{c}$ lies on the physical strip instead, the correct expressions for the minimal K_2 -matrices can be obtained by the interchange $c \leftrightarrow -c$, $-\circ- \leftrightarrow -\circ-$ (together with $d \leftrightarrow -d$ for $SU(N)$). (Note that this leaves $c-\circ-$ unchanged.) In section 4 we shall add CDD factors making the PCM boundary S -matrices invariant under this interchange.

3.4.1 $SU(N)$

The minimal boundary S -matrices for $SU(N)$ are

$$K_1(\theta) = \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2}\right)} \text{---}\bigcirc \quad \text{with } \bigcirc = 0 \quad (3.21)$$

where Γ is the gamma function, and

$$K_2(\theta) = \frac{-1}{(1-c\theta)} \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{-\theta}{2i\pi}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2i\pi d}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{\theta}{2i\pi}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2i\pi d}\right)} (\text{---} + c\theta \text{---}\bigcirc) \quad (3.22)$$

with $\bigcirc = \frac{ih}{\pi}(\frac{1}{c} + \frac{1}{d})$.

(Note that in the limit $c, d \rightarrow \infty$, with $\frac{cd}{c+d}\bigcirc$ fixed, $K_2(\theta) \rightarrow K_1(\theta)$, as we would expect.)

3.4.2 $SO(N)$

There are two types of minimal boundary S -matrix, corresponding to E symmetric or antisymmetric, for each of the $K_1(\theta)$ and $K_2(\theta)$ forms in the case $G = SO(N)$. (The symmetric $K_2(\theta)$ type was investigated in [8] and, for $M = 1$, in [35]. The $K_2(\theta)$ type with antisymmetric E has been considered in [36].) The minimal boundary S -matrices of the $K_1(\theta)$ form are

$$K_1(\theta) = \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{4} + \frac{1}{2h}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{4} + \frac{1}{2h}\right)} \text{---}\bigcirc \quad (3.23)$$

with $(\text{---}\bigcirc)^t = \text{---}\bigcirc$, $\bigcirc = 0$, and

$$K_1(\theta) = \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{3}{4} + \frac{1}{2h}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{3}{4} + \frac{1}{2h}\right)} \text{---}\bigcirc \quad (3.24)$$

with $(\text{---}\bigcirc)^t = -(\text{---}\bigcirc)$, $(\Rightarrow \bigcirc = 0)$. Those of the $K_2(\theta)$ form are

$$K_2(\theta) = \frac{-1}{(1-c\theta)} \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{-\theta}{2i\pi}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{4} + \frac{1}{2h}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{\theta}{2i\pi}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{4} + \frac{1}{2h}\right)} \times \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2i\pi c}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2i\pi c}\right)} (\text{---} + c\theta \text{---}\bigcirc) \quad (3.25)$$

with $(\text{---}\bigcirc)^t = \text{---}\bigcirc$, $c\bigcirc = \frac{2ih}{\pi}$, and

$$K_2(\theta) = \frac{-1}{(1-c\theta)} \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{-\theta}{2i\pi}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{3}{4} + \frac{1}{2h}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{\theta}{2i\pi}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{3}{4} + \frac{1}{2h}\right)}$$

$$\times \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2i\pi c}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2i\pi c}\right)} (- + c\theta \text{---} \text{---}) \quad (3.26)$$

with $(\text{---})^t = -(\text{---})$, $(\Rightarrow \bigcirc = 0)$.

(Note that in the limit $c \rightarrow \infty$, with $c\bigcirc$ fixed, the symmetric (respectively antisymmetric) $K_2(\theta)$ tends to the symmetric (respectively antisymmetric) $K_1(\theta)$, again as expected.)

3.4.3 $Sp(N)$

For $Sp(N)$ there are again two minimal solutions of the form $K_1(\theta)$:

$$K_1(\theta) = \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{4} + \frac{1}{2h}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{4} + \frac{1}{2h}\right)} \text{---} \quad (3.27)$$

with $\rightarrow(\text{---})^t \rightarrow = \text{---}$, $(\Rightarrow \bigcirc = 0)$, and

$$K_1(\theta) = \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{3}{4} + \frac{1}{2h}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{3}{4} + \frac{1}{2h}\right)} \text{---} \quad (3.28)$$

with $\rightarrow(\text{---})^t \rightarrow = -(\text{---})$, $\bigcirc = 0$. The two minimal boundary S -matrices of the form $K_2(\theta)$ are

$$K_2(\theta) = \frac{-1}{(1 - c\theta)} \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{-\theta}{2i\pi}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{4} + \frac{1}{2h}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{\theta}{2i\pi}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{4} + \frac{1}{2h}\right)} \times \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2i\pi c}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2i\pi c}\right)} (- + c\theta \text{---} \text{---}) \quad (3.29)$$

with $\rightarrow(\text{---})^t \rightarrow = \text{---}$, $(\Rightarrow \bigcirc = 0)$, and

$$K_2(\theta) = \frac{-1}{(1 - c\theta)} \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{-\theta}{2i\pi}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{3}{4} + \frac{1}{2h}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{3}{4}\right) \Gamma\left(\frac{\theta}{2i\pi}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{3}{4} + \frac{1}{2h}\right)} \times \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2i\pi c}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2i\pi c}\right)} (- + c\theta \text{---} \text{---}) \quad (3.30)$$

with $\rightarrow(\text{---})^t \rightarrow = -(\text{---})$, $c\bigcirc = \frac{2ih}{\pi}$.

(As $c \rightarrow \infty$, with $c\bigcirc$ fixed, the $K_2(\theta)$ with $\rightarrow(\text{---})^t \rightarrow = \pm(\text{---})$ tends to the $K_1(\theta)$ with the respective property, as expected.)

3.4.4 $SU(N)$ -conjugating

Here it is only $K_1(\theta)$ that provides valid boundary S -matrices. There are two minimal possibilities, with symmetric and antisymmetric $\neg\circ$,

$$K_1(\theta) = \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{4}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{4} + \frac{1}{2h}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{4}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{4} + \frac{1}{2h}\right)} \neg\circ \quad \text{with } (\neg\circ)^t = \neg\circ, \quad \text{and} \quad (3.31)$$

$$K_1(\theta) = \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{4}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{3}{4} + \frac{1}{2h}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{4}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{3}{4} + \frac{1}{2h}\right)} \neg\circ \quad \text{with } (\neg\circ)^t = -(\neg\circ). \quad (3.32)$$

We have been unable to make contact between our solutions and a rational limit of the trigonometric solutions in [11].

3.5 Constraints on E : the symmetric-space correspondence

We now turn our attention to the constraints imposed on the matrices E . We recall that

$$E^\dagger = E \quad \text{and} \quad E^2 = I \quad (3.33)$$

were to be imposed in all cases, except that of $SU(N)$ -conjugating, due to unitarity and analyticity. For the case of $SU(N)$ -conjugating (3.33) is replaced by

$$E^\dagger E = I \quad \text{and} \quad \det E = 1. \quad (3.34)$$

The further constraints particular to the different groups were

Group	$K_1(\theta)$	$K_2(\theta)$
$SU(N)$	$\text{Tr}(E) = 0$	$\text{Tr}(E) = \frac{ih}{\pi} \left(\frac{1}{c} + \frac{1}{d} \right)$
$SO(N)$	$\text{Tr}(E) = 0,$ $E^t = \pm E$	$\text{Tr}(E) = \frac{2ih}{c\pi}, \quad \text{or} \quad \text{Tr}(E) = 0,$ $E^t = E \quad \quad \quad E^t = -E$
$Sp(2n)$	$\text{Tr}(E) = 0,$ $JE^t J = \pm E$	$\text{Tr}(E) = 0, \quad \text{or} \quad \text{Tr}(E) = \frac{2ih}{c\pi},$ $JE^t J = E \quad \quad \quad JE^t J = -E$
$SU(N)$ - conjugating	$E^t = \pm E$	no solution

3.5.1 $SU(N)$

We begin by considering $SU(N)$, where in addition to the constraints imposed by unitarity and analyticity we have a single extra constraint on the trace of E . From the first two

constraints we can express E as the conjugate of a diagonal matrix X by an $SU(N)$ matrix,

$$E = Q^\dagger X Q \quad \text{with } Q \in SU(N)$$

where X is of the form

$$X = \begin{pmatrix} I_M & 0 \\ 0 & -I_{N-M} \end{pmatrix},$$

I_M is the $M \times M$ identity matrix and $0 \leq M \leq N$ (see appendix 5.2.1). By the cyclicity of trace, if we impose the trace condition for K_1 then N must be even and equal to $2M$. If we impose the condition for K_2 we obtain a condition on c and d , and so find

$K_1(\theta)$	$K_2(\theta)$
$E = Q_1^\dagger \begin{pmatrix} I_{N/2} & 0 \\ 0 & -I_{N/2} \end{pmatrix} Q_1$	$E = Q_2^\dagger \begin{pmatrix} I_M & 0 \\ 0 & -I_{N-M} \end{pmatrix} Q_2 \quad \text{where } \frac{1}{c} + \frac{1}{d} = \frac{\pi(2M-N)}{i\hbar}$

with $Q_i \in SU(N)$. We can see that the case K_1 corresponds to the limit of K_2 in which we take $M = \frac{N}{2}$, that is the $c, d \rightarrow \infty$, as we would expect.

Thus we have parametrized the possibilities for E with a matrix $Q \in SU(N)$ and an integer M . Once M is fixed, the suitable E form a space isomorphic to the symmetric space

$$\frac{SU(N)}{S(U(M) \times U(N-M))}$$

where the correspondence is between an element $E = Q^\dagger X Q$ and the left coset of $H = S(U(M) \times U(N-M))$ by Q . In the same way the possible E for $K_1(\theta)$ form a space isomorphic to

$$\frac{SU(N)}{S(U(N/2) \times U(N/2))} \quad (\text{only for } N \text{ even}).$$

3.5.2 $SO(N)$

We now consider $SO(N)$, where in addition to the constraints associated with $SU(N)$ we also have $E^t = \pm E$. We consider first the case $E^t = E$:

$$E^\dagger = E, \quad E^t = E \Rightarrow E^* = E.$$

So E is a symmetric real matrix, and we can diagonalize it by conjugating with a matrix $R \in SO(N)$. Since E squares to the identity the diagonal matrix must be of the form

$$X = \begin{pmatrix} I_M & 0 \\ 0 & -I_{N-M} \end{pmatrix}.$$

Then imposing the constraints on the trace of E is the same as for $SU(N)$ and we have

$K_1(\theta)$	$K_2(\theta)$
$E = R_1^t \begin{pmatrix} I_{N/2} & 0 \\ 0 & -I_{N/2} \end{pmatrix} R_1$	$E = R_2^t \begin{pmatrix} I_M & 0 \\ 0 & -I_{N-M} \end{pmatrix} R_2$ where $c = \frac{2ih}{\pi(2M-N)}$

with $R_i \in SO(N)$. In a similar way to the $SU(N)$ case, once M is fixed, the space of matrices E is isomorphic to

$$\frac{SO(N)}{S(O(M) \times O(N-M))},$$

with the E for $K_1(\theta)$ isomorphic to

$$\frac{SO(N)}{S(O(N/2) \times O(N/2))}.$$

Thus, in the same way as for $SU(N)$, we have an isomorphism between the space of allowed E and the symmetric spaces (see appendix 5.2.2 for more details).

The remaining case to consider is that of antisymmetric E . We find in this case (see appendix 5.2.6) that the matrices E form a space isomorphic to two copies of the symmetric space

$$\frac{SO(N)}{U(N/2)}.$$

3.5.3 $Sp(N)$

In addition to the $SU(N)$ constraints, for $Sp(N)$ we also have $JE^t J = \pm E$. We consider first the case $JE^t J = -E$, with

$$E^2 = I \quad \text{and} \quad JE^t J = -E \Rightarrow JE^t J E = -I \Rightarrow E^t J E = J$$

so that, since we also know $E \in U(N)$, we must have $E \in Sp(N)$. After appealing to an argument involving quaternionic matrices (appendix 5.2.3) we find that the space of allowed E for K_2 is isomorphic to

$$\frac{Sp(N)}{Sp(M) \times Sp(N-M)}.$$

In the case of $K_1(\theta)$ we again require $M = \frac{N}{2}$ and the E -space is isomorphic to

$$\frac{Sp(N)}{Sp(N/2) \times Sp(N/2)}.$$

For $K_1(\theta)$ with $JE^tJ = +E$, E is conjugate over $\mathbb{C}$ to

$$\begin{pmatrix} I_{N/2} & 0 \\ 0 & -I_{N/2} \end{pmatrix}$$

as $Tr(E) = 0$. We find (see appendix 5.2.7) that the allowed E form a space isomorphic to

$$\frac{Sp(N)}{U(N/2)}.$$

3.5.4 $SU(N)$ -conjugating

The last case to consider is that of $SU(N)$ -conjugating. We recall that in this case the constraints due to unitarity and analyticity were slightly modified, to

$$E^\dagger E = I \quad \text{and} \quad \det E = 1.$$

Crossing-unitarity imposed $E^t = \pm E$. Taking the symmetric case first, the allowed E form a set

$$\{E | E^\dagger E = I, E^t = E, \det E = 1\}$$

which turns out (see appendix 5.2.4) to be isomorphic to

$$\frac{SU(N)}{SO(N)}.$$

Lastly, we turn to the antisymmetric case

$$\{E | E^\dagger E = I, E^t = -E, \det E = 1\}.$$

This we find (appendix 5.2.5) is isomorphic to

$$\{1, \omega^2\} \times \frac{SU(N)}{Sp(N)} \quad \text{where any } \omega \text{ s.t. } \omega^N = -1 \text{ is chosen.}$$

4 The PCM boundary S -matrices

In this section we construct the boundary S -matrices for the principal chiral model. We recall [29] that the bulk model S -matrix has $G \times G$ symmetry and is constructed as

$$S_{PCM}(\theta) = X_{11}(\theta) \left(S_L(\theta) \otimes S_R(\theta) \right) \quad (4.1)$$

where $X_{11}(\theta)$ is the CDD factor for the PCM and $S_{L,R}(\theta)$ are left and right copies of the minimal S -matrix possessing G -symmetry. Following this prescription, we shall use the minimal K -matrices from the previous section to construct boundary S -matrices for the PCM on a half-line. We then go on to explore their symmetries and make connection with the classical results of section two.

4.1 The CDD factors

Introducing the CDD factor, $X_{11}(\theta)$, into the bulk S -matrix for the PCM requires that we introduce an extra factor, $Y_{11}(\theta)$ (or $Y_{1\bar{1}}(\theta)$ in the case of $SU(N)$ conjugating), into the boundary S -matrix in order to satisfy crossing-unitarity. We construct the boundary S -matrix for the PCM as

$$K_{PCM}(\theta) = Y_{11}(\theta) \left(K_L(\theta) \otimes K_R(\theta) \right) \quad (4.2)$$

where $K_{L,R}(\theta)$ are left and right copies of the same type of minimal K -matrix, chosen from among the possibilities classified in section three. In order that $K_{PCM}(\theta)$ satisfy the crossing-unitarity equation with $S_{PCM}(\theta)$ we require

$$\begin{aligned} Y_{11}(i\pi - \theta)Y_{11}(i\pi + \theta) &= 1 & SU(N), \\ \frac{Y_{11}(\frac{i\pi}{2} - \theta)}{Y_{11}(\frac{i\pi}{2} + \theta)} &= X_{11}(2\theta) & SO(N) \text{ and } Sp(N), \\ \frac{Y_{1\bar{1}}(\frac{i\pi}{2} - \theta)}{Y_{1\bar{1}}(\frac{i\pi}{2} + \theta)} &= X_{11}(2\theta) & SU(N) - \text{conjugating}. \end{aligned} \quad (4.3)$$

The CDD factors for the bulk PCM S -matrices are

$$\begin{aligned} X_{11}(\theta) &= (2)_\theta = X_{1\bar{1}}(i\pi - \theta) & SU(N), \\ X_{11}(\theta) &= (2)_\theta (h - 2)_\theta & SO(N) \text{ and } Sp(N), \end{aligned} \quad (4.4)$$

$$\text{where } (x)_\theta = \frac{\sinh(\frac{\theta}{2} + \frac{i\pi x}{2h})}{\sinh(\frac{\theta}{2} - \frac{i\pi x}{2h})}.$$

We find [30] the following candidates for the Y functions:

$$\begin{aligned} Y_{11}(\theta) &= -(1 - h)_\theta & SU(N) \\ Y_{11}(\theta) &= -\left(\frac{h}{2} + 2\right)_\theta \left(\frac{h}{2} + 1\right)_\theta (1 - h)_\theta & SO(N) \text{ and } Sp(N) \\ Y_{1\bar{1}}(\theta) &= \left(\frac{h}{2} + 2\right)_\theta \left(\frac{h}{2} + 1\right)_\theta & SU(N) - \text{conjugating}, \end{aligned} \quad (4.5)$$

and note that none of these factors have poles on the physical strip.

We still have freedom to multiply by an arbitrary boundary CDD factor. That is, we can replace any of the above $Y(\theta)$ factors by $g(\theta)Y(\theta)$, where $g(\theta)$ is a CDD factor. This allows us to introduce simple poles into the PCM boundary S -matrices. In the case where we construct a PCM S -matrix using left and right copies of $K_2(\theta)$, we wish to introduce the CDD factor⁸

$$\begin{aligned} g(\theta) &= \left(\frac{h}{ci\pi} \right)_\theta \left(h - \frac{h}{di\pi} \right)_\theta \quad SU(N), \\ g(\theta) &= \left(\frac{h}{ci\pi} \right)_\theta \left(h - \frac{h}{ci\pi} \right)_\theta \quad SO(N) \text{ and } Sp(N) \end{aligned} \quad (4.6)$$

which gives a simple pole at $\theta = \frac{1}{c}$, corresponding to the formation of a boundary bound state.

4.2 The boundary S -matrices

We now list the full PCM boundary S -matrices for the various G . We make use of the relation

$$\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin(\pi z)}.$$

We will also require the scalar factors

$$\begin{aligned} \eta(\theta) &= \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2}\right)}, \quad \nu(\theta) = \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{-\theta}{2i\pi}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2h}\right) \Gamma\left(\frac{\theta}{2i\pi}\right)}, \\ \mu(\theta) &= \frac{1}{4\pi^2 c^2} \frac{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{1}{2} - \frac{1}{2i\pi c}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{\theta}{2i\pi} - \frac{1}{2i\pi c}\right)}{\Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2} - \frac{1}{2i\pi c}\right) \Gamma\left(\frac{-\theta}{2i\pi} + 1 + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{-\theta}{2i\pi} + 1 - \frac{1}{2i\pi c}\right)}, \\ \lambda(\theta) &= \frac{1}{4\pi^2 c^2} \frac{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{M}{N} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{-\theta}{2i\pi} + 1 - \frac{M}{N} - \frac{1}{2i\pi c}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{\theta}{2i\pi} - \frac{1}{2i\pi c}\right)}{\Gamma\left(\frac{\theta}{2i\pi} + \frac{M}{N} + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{\theta}{2i\pi} + 1 - \frac{M}{N} - \frac{1}{2i\pi c}\right) \Gamma\left(\frac{-\theta}{2i\pi} + 1 + \frac{1}{2i\pi c}\right) \Gamma\left(\frac{-\theta}{2i\pi} + 1 - \frac{1}{2i\pi c}\right)}, \\ \text{and} \quad \epsilon_{n,m}(\theta) &= \frac{\Gamma\left(\frac{\theta}{2i\pi} + \frac{n}{4}\right) \Gamma\left(\frac{-\theta}{2i\pi} + \frac{m}{4} + \frac{1}{2h}\right)}{\Gamma\left(\frac{-\theta}{2i\pi} + \frac{n}{4}\right) \Gamma\left(\frac{\theta}{2i\pi} + \frac{m}{4} + \frac{1}{2h}\right)}. \end{aligned}$$

The PCM S -matrices can then be written as follows.

⁸We are assuming, as in section 3.4, that $M \leq \frac{N}{2}$ so that $\frac{1}{c}$ is on the physical strip. As stated in section 3.4, the resulting PCM boundary S -matrix will possess a $c \leftrightarrow -c$ (and $d \leftrightarrow -d$ for $G = SU(N)$) symmetry and so will be correct for $M \geq \frac{N}{2}$ also.

4.2.1 $SU(N)$

We have found two types of boundary S -matrix for $SU(N)$,

$$K_{PCM}(\theta) = -(1-h)_\theta \mu(\theta) \left(\nu(\theta)(I + c\theta E_L) \otimes \nu(\theta)(I + c\theta E_R) \right) \quad (4.7)$$

(whose $c \rightarrow \infty$ limit:

$$K_{PCM}(\theta) = -(1-h)_\theta \left(\eta(\theta) E_L \otimes \eta(\theta) E_R \right) \quad (4.8)$$

is a valid PCM boundary scattering matrix) where⁹

$$E_{L/R} \in \frac{SU(N)}{S(U(N/2) \times U(N/2))},$$

and

$$K_{PCM}(\theta) = -(1-h)_\theta \lambda(\theta) \left(\nu(\theta)(I + c\theta E_L) \otimes \nu(\theta)(I + c\theta E_R) \right) \quad (4.9)$$

where

$$E_{L/R} \in \frac{SU(N)}{S(U(N-M) \times U(M))}.$$

4.2.2 $SO(N)$

For $SO(N)$ three types have been found,

$$K_{PCM}(\theta) = - \left(\frac{h}{2} + 2 \right)_\theta \left(\frac{h}{2} + 1 \right)_\theta (1-h)_\theta \mu(\theta) \left(\nu(\theta) \epsilon_{3,3}(\theta)(I + c\theta E_L) \otimes \nu(\theta) \epsilon_{3,3}(\theta)(I + c\theta E_R) \right) \quad (4.10)$$

(whose $c \rightarrow \infty$ limit:

$$K_{PCM}(\theta) = - \left(\frac{h}{2} + 2 \right)_\theta \left(\frac{h}{2} + 1 \right)_\theta (1-h)_\theta \left(\eta(\theta) \epsilon_{3,3}(\theta) E_L \otimes \eta(\theta) \epsilon_{3,3}(\theta) E_R \right) \quad (4.11)$$

is a valid PCM boundary scattering matrix) where¹⁰

$$E_{L/R} \in \frac{SO(N)}{U(N/2)} \times \{+1, -1\},$$

$$K_{PCM}(\theta) = - \left(\frac{h}{2} + 2 \right)_\theta \left(\frac{h}{2} + 1 \right)_\theta (1-h)_\theta \left(\eta(\theta) \epsilon_{3,1}(\theta) E_L \otimes \eta(\theta) \epsilon_{3,1}(\theta) E_R \right) \quad (4.12)$$

⁹We are using the symmetric space notation G/H here to denote the relevant translated Cartan immersion. Details are given in appendix 5.

¹⁰The factor $\{+1, -1\}$ indicates that the space containing $E_{L/R}$ is a twofold copy of the symmetric space – no group structure is implied. See appendix 5.2.6 for details.

where

$$E_{L/R} \in \frac{SO(N)}{S(O(N/2) \times O(N/2))},$$

and

$$K_{PCM}(\theta) = - \left(\frac{h}{2} + 2 \right)_\theta \left(\frac{h}{2} + 1 \right)_\theta (1-h)_\theta \mu(\theta) \left(\nu(\theta) \epsilon_{3,1}(\theta) (I + c\theta E_L) \otimes \nu(\theta) \epsilon_{3,1}(\theta) (I + c\theta E_R) \right) \quad (4.13)$$

where

$$E_{L/R} \in \frac{SO(N)}{S(O(N-M) \times O(M))} \quad \text{and } c = \frac{2ih}{\pi(2M-N)} \text{ in } \mu(\theta).$$

4.2.3 $Sp(N)$

Three types of $K_{PCM}(\theta)$ have also been found for $Sp(N)$,

$$K_{PCM}(\theta) = - \left(\frac{h}{2} + 2 \right)_\theta \left(\frac{h}{2} + 1 \right)_\theta (1-h)_\theta \mu(\theta) \left(\nu(\theta) \epsilon_{3,1}(\theta) (I + c\theta E_L) \otimes \nu(\theta) \epsilon_{3,1}(\theta) (I + c\theta E_R) \right) \quad (4.14)$$

(whose $c \rightarrow \infty$ limit:

$$K_{PCM}(\theta) = - \left(\frac{h}{2} + 2 \right)_\theta \left(\frac{h}{2} + 1 \right)_\theta (1-h)_\theta \left(\eta(\theta) \epsilon_{3,1}(\theta) E_L \otimes \eta(\theta) \epsilon_{3,1}(\theta) E_R \right) \quad (4.15)$$

is a valid PCM boundary scattering matrix) where

$$E_{L/R} \in \frac{Sp(N)}{U(N/2)},$$

$$K_{PCM}(\theta) = - \left(\frac{h}{2} + 2 \right)_\theta \left(\frac{h}{2} + 1 \right)_\theta (1-h)_\theta \left(\eta(\theta) \epsilon_{3,3}(\theta) E_L \otimes \eta(\theta) \epsilon_{3,3}(\theta) E_R \right) \quad (4.16)$$

where

$$E_{L/R} \in \frac{Sp(N)}{Sp(N/2) \times Sp(N/2)},$$

and

$$K_{PCM}(\theta) = - \left(\frac{h}{2} + 2 \right)_\theta \left(\frac{h}{2} + 1 \right)_\theta (1-h)_\theta \mu(\theta) \left(\nu(\theta) \epsilon_{3,3}(\theta) (I + c\theta E_L) \otimes \nu(\theta) \epsilon_{3,3}(\theta) (I + c\theta E_R) \right) \quad (4.17)$$

where

$$E_{L/R} \in \frac{Sp(N)}{Sp(N-M) \times Sp(M)} \quad \text{and } c = \frac{2ih}{\pi(2M-N)} \text{ in } \mu(\theta).$$

4.2.4 $SU(N)$ -conjugating

Lastly, we have found two types of representation-conjugating boundary S -matrix for $SU(N)$

$$K_{PCM}(\theta) = \left(\frac{h}{2} + 2\right)_\theta \left(\frac{h}{2} + 1\right)_\theta \left(\epsilon_{1,1}(\theta)E_L \otimes \epsilon_{1,1}(\theta)E_R\right) \quad (4.18)$$

where

$$E_{L/R} \in \frac{SU(N)}{SO(N)},$$

and

$$K_{PCM}(\theta) = \left(\frac{h}{2} + 2\right)_\theta \left(\frac{h}{2} + 1\right)_\theta \left(\epsilon_{1,3}(\theta)E_L \otimes \epsilon_{1,3}(\theta)E_R\right) \quad (4.19)$$

where

$$E_{L/R} \in \{1, \omega^2\} \times \frac{SU(N)}{Sp(N)} \quad (\omega^N = -1).$$

4.3 Symmetries of the PCM boundary S -matrices

We now consider the symmetries possessed by the PCM boundary S -matrices. Before looking at the surviving group symmetries at the boundary, we first point out a symmetry possessed by those S -matrices constructed from the $K_2(\theta)$ -type minimal solution.

4.3.1 $M \leftrightarrow N - M$ symmetry

This was first noted, for $SU(N)$ diagonal boundary scattering, in [33]. The Grassmannian symmetric spaces

$$\frac{SU(N)}{S(U(M) \times U(N - M))} \quad \frac{SO(N)}{S(O(M) \times O(N - M))} \quad \frac{Sp(N)}{Sp(M) \times Sp(N - M)}$$

are all invariant under $M \leftrightarrow N - M$. Consequently, we might expect that the PCM boundary S -matrices constructed using matrices $E_{L/R}$ lying in translated Cartan constructions of these symmetric spaces would also respect this symmetry. This is exactly what we find for the $K_{PCM}(\theta)$ matrices (4.9), (4.13) and (4.17).

To see this invariance, we consider how the exchange $M \leftrightarrow N - M$ affects the degrees of freedom in the K -matrices. The matrices $E_{L/R}$ are constructed as

$$E_{L/R} = U_{L/R} X U_{L/R}^{-1} \quad \text{where} \quad U_{L/R} \in SU(N), SO(N) \text{ or } Sp(N) \quad \text{and} \quad (4.20)$$

$$X = \left(\begin{array}{c|c} I_M & 0 \\ \hline 0 & -I_{N-M} \end{array} \right), \quad \text{so under } M \leftrightarrow N-M \quad X \mapsto \check{X} = \left(\begin{array}{c|c} I_{N-M} & 0 \\ \hline 0 & -I_M \end{array} \right).$$

Thus under $M \leftrightarrow N-M$, $E_{L/R} \mapsto \check{E}_{L/R} = U_{L/R} \check{X} U_{L/R}^{-1}$. Taking traces we see that $c \leftrightarrow -c$ (and $d \leftrightarrow -d$) under the exchange. Now note that the scalar factor $\mu(\theta)$ is invariant under $c \leftrightarrow -c$ and $\lambda(\theta)$ is invariant under $c \leftrightarrow -c$, $M \leftrightarrow N-M$. So $K_{PCM}(\theta) \mapsto \check{K}_{PCM}(\theta)$ where

$$(I + c\theta E_{L/R}) \mapsto (I - c\theta \check{E}_{L/R}).$$

Now

$$-\check{E}_{L/R} = U_{L/R}(-\check{X})U_{L/R}^{-1} = U_{L/R}OXO^{-1}U_{L/R}^{-1}$$

where

$$O = \left(\begin{array}{c|c} 0 & I_{N-M} \\ \hline \pm I_M & 0 \end{array} \right).$$

We choose the sign $\pm$ to ensure that $\det O = 1$ and (noting that when N and M are even $O \in Sp(N)$) we see that $U_{L/R} \in G = SU(N)$, $SO(N)$ or $Sp(N) \implies (U_{L/R}O) \in G$.

Thus, if we denote by $K_{PCM}(\theta; U_L, U_R)$ the matrix constructed using

$$E_{L/R} = U_{L/R} X U_{L/R}^{-1} \tag{4.21}$$

and by $\check{K}_{PCM}(\theta; U_L, U_R)$ the image of this under the exchange $M \leftrightarrow N-M$, we have

$$\check{K}_{PCM}(\theta; U_L, U_R) = K_{PCM}(\theta; U_L O, U_R O). \tag{4.22}$$

So we see that the $K_{PCM}(\theta)$ matrices do respect this invariance of the symmetric spaces, in the sense that the action of the exchange $M \leftrightarrow N-M$ on the K -matrices is simply a translation in the parameter space.

A further consequence of this emerges if we consider the pole structure of these K -matrices. Restricting to the choice of parameter $c = d = \frac{2iN}{\pi(2M-N)}$ in the case $G = SU(N)$, so that it is analogous to the $G = SO(N)$, $Sp(N)$ cases, there is exactly one simple pole on the physical strip at either $\theta = \frac{1}{c}$ or $\theta = -\frac{1}{c}$ (since $M \neq \frac{N}{2}$). If we interpret the simple pole as the formation of a boundary bound state at this rapidity, then the bound state is in a representation projected onto by either $P_L^+ \otimes P_R^+$ or $P_L^- \otimes P_R^-$, respectively, where

$$P_{L/R}^\pm = \frac{1}{2}(I \pm E_{L/R}) = U_{L/R} \left(\frac{1}{2}(I \pm X) \right) U_{L/R}^{-1} \tag{4.23}$$

We note

$$\frac{1}{2}(I + X) = \left(\begin{array}{c|c} I_M & 0 \\ \hline 0 & 0 \end{array} \right) \quad \text{and} \quad \frac{1}{2}(I - X) = \left(\begin{array}{c|c} 0 & 0 \\ \hline 0 & I_{N-M} \end{array} \right).$$

We find that $\pm \frac{1}{c}$ lies on the physical strip as $M \lesssim \frac{N}{2}$, and so the boundary bound state representation is always the smaller of the two projection spaces. We plan to investigate further the spectrum of boundary bound states in future work, but for the moment we return to consider the surviving remnant of group symmetry at the boundary and make connections with the classical boundary conditions of section two.

4.3.2 Boundary group symmetry in the non-conjugating cases

We recall [29] that the principal chiral model in the bulk possesses a global $G \times G$ symmetry, respected by the bulk S -matrices. In section two we saw that the introduction of a boundary in the classical PCM generally breaks the $G \times G$ symmetry, so that only a remnant survives, the nature of which is dictated by the boundary condition. In particular we saw in section 2.2.1 that the boundary condition (2.11),

$$g(0) \in k_L H k_R^{-1} \quad \text{where } k_{L/R} \text{ parametrize left/right cosets of } H \in G,$$

preserves $k_L H k_L^{-1} \times k_R H k_R^{-1}$.

Turning our attention to the PCM boundary S -matrices, we find that $K_{PCM}(\theta; k_L, k_R)$ is invariant under exactly this symmetry. That is,

$$[K_{PCM}(\theta; k_L, k_R), k_L H k_L^{-1} \times k_R H k_R^{-1}] = 0. \quad (4.24)$$

We begin with the Grassmannian cases, where it is enough to show that

$$[(I + c\theta E_L) \otimes (I + c\theta E_R), k_L h_L k_L^{-1} \times k_R h_R k_R^{-1}] = 0,$$

where (for subscripts L and R) h are arbitrary elements of H , $E = kXk^{-1}$ and $k \in G = SU(N)$, $SO(N)$ or $Sp(N)$. But this is immediate: $Xh = hX$, since H is constructed to be precisely those elements in G which commute with X .

For the case $G/H = SO(2n)/U(n)$ (respectively $G/H = Sp(2n)/U(n)$) we note (appendix 5.2.6, resp. 5.2.7) that $E = ikJk^{-1}$ where $k \in G$, and that $H = U(n)$ is constructed as those elements in G satisfying $Jh = hJ$, giving the required result.

4.3.3 Boundary group symmetry in the $SU(N)$ -conjugating case

The cases of $SU(N)/SO(N)$ and $SU(N)/Sp(N)$ are a little more subtle. Performing similar calculations to the above (and again leaving the L/R suffix implicit) we find, on

constructing E as in appendices 5.2.4, 5.2.5, that $E k h k^{-1} = (k h k^{-1})^* E$, which implies

$$K_{PCM}(\theta; k_L, k_R) (k_L H_L k_L^{-1} \times k_R H_R k_R^{-1}) = (k_L H_L k_L^{-1} \times k_R H_R k_R^{-1})^* K_{PCM}(\theta; k_L, k_R). \quad (4.25)$$

Such a result is not surprising, since in this case $K_{PCM}(\theta) : V_L \otimes V_R \rightarrow \bar{V}_L \otimes \bar{V}_R$. It is straightforward to obtain a symmetry relation in which, as earlier in section 3.2.2, we consider a boundary S -matrix which is an endomorphism of $(V_L \oplus \bar{V}_L) \otimes (V_R \oplus \bar{V}_R)$. We do not give details.

4.4 Concluding summary

When G/H is a symmetric space, the classical boundary condition $g(0) \in k_L H k_R^{-1}$ preserves the local PCM conserved charges necessary for integrability. Thus, as stated in section 2.2.1, the possible BCs are parametrized by a moduli space $G/H \times G/H^{11}$. We have also found boundary S -matrices which are parametrized by $G/H \times G/H$. Further, we find that the global symmetry which survives in the presence of this BC is precisely that which commutes with the boundary S -matrix $K_{PCM}(\theta; k_L, k_R)^{12}$. So we finish with this

Claim: The principal chiral model on G is classically integrable with boundary condition $g(0) \in k_L H k_R^{-1}$, where $k_{L/R} \in G$ and G/H is a symmetric space; and it remains integrable at the quantum level, where its boundary S -matrix is $K_{PCM}(\theta; k_L, k_R)$.

Additional comment for v3

Recently a paper [34] has appeared which deals with $osp(m|n)$ spin chain models. Its results are related to ours for the $G = SO(N)$ and $G = Sp(N)$ cases.

Acknowledgments

We should like to thank Tony Sudbery and Ian McIntosh for discussions of symmetric spaces. Our thanks also go to Gérard Watts for pointing out an error in our original discussion of crossing unitarity. NJM would like to thank Patrick Dorey and Ed Corrigan for helpful discussions, and Bernard Piette, Paul Fendley and Jonathan Evans for email exchanges. BJS would like to thank Gustav Delius, Brett Gibson and Mark Kambites for discussions. Finally NJM thanks the Centre de Recherches Mathématiques, U de Montréal, where this work was begun during the ‘Quantum Integrability 2000’ program, for hospitality and financial support, and BJS thanks the UK EPSRC for a D.Phil. studentship.

¹¹up to a discrete ambiguity, as further explained in appendix 6.1

¹²with the subtlety noted above in the case of $SU(N)$ -conjugated

Finally, we should like to thank Gérard Watts for pointing out an error (in our use of the crossing-unitarity relation) in earlier versions which has led to the changes (namely the slightly greater freedom in some scalar prefactors) in v3.

5 Appendix: Symmetric spaces and the Cartan immersion

Under the action of an involutive automorphism α (which may or may not be inner), a Lie algebra splits into eigenspaces $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{k}$ of eigenvalue $+1$ ($\mathfrak{h}$) and -1 ($\mathfrak{k}$), with

$$[\mathfrak{h}, \mathfrak{h}] \subset \mathfrak{h}, \quad [\mathfrak{h}, \mathfrak{k}] \subset \mathfrak{k}, \quad [\mathfrak{k}, \mathfrak{k}] \subset \mathfrak{h}.$$

The subgroup H generated by $\mathfrak{h}$ is compact, and we have taken G to be compact (type I) rather than maximally non-compact (type III). For the classical groups these are the groups $G = SU(N), SO(N)$ and $Sp(N)$ (where the argument of Sp is understood always to be even) themselves, along with those described in the table below. The dimension is $\dim G - \dim H$, and the automorphism is given by its action on U in the defining representation, where X is the diagonal matrix with M $+1$ s and $N - M$ -1 s and J is the symplectic form matrix, which is block-diagonal with $N/2$ blocks $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and satisfies $J^2 = -I_N$.

symmetric space	dimension	automorphism
$SU(N)/S(U(N-M) \times U(M))$	$2M(N-M)$	$U \mapsto XUX$
$SO(N)/SO(N-M) \times SO(M)$	$M(N-M)$	$U \mapsto XUX$
$Sp(N)/Sp(N-M) \times Sp(M)$	$M(N-M)$	$U \mapsto XUX$
$SU(N)/SO(N)$	$\frac{N(N+1)}{2} - 1$	$U \mapsto U^*$
$SU(N)/Sp(N)$	$\frac{N(N-1)}{2} - 1$	$U \mapsto -JU^*J$
$SO(2n)/U(n)$	$n(n-1)$	$U \mapsto -JUJ$
$Sp(2n)/U(n)$	$n(n+1)$	$U \mapsto -JUJ$

(We refer to the first three as the ‘Grassmannian’ cases.)

5.1 The Cartan immersion

The Cartan immersion constructs G/H as a subspace of G (due to Cartan, and described briefly in [17] or more fully in [18] (vol.II, sect.10, prop.4). Lifting α in the natural way

from the algebra to the group (so that $\alpha(h) = h$ for all $h \in H$), under

$$\begin{aligned} gH &\mapsto \alpha(g)g^{-1} \\ \text{we have } G/H &= \{\alpha(g)g^{-1} | g \in G\}. \end{aligned}$$

(This statement, of course, depends crucially on the fact that we have chosen H so that it consists of *all* elements of G invariant under α ; for the more general case see [39, 40].)

We then have $\alpha(k) = k^{-1}$ for all $k \in \frac{G}{H} \hookrightarrow G$. Defining

$$\mathcal{M} = \{k \in G | \alpha(k) = k^{-1}\}, \quad (5.1)$$

it turns out [39, 40] that G/H is in 1-1 correspondence with $\mathcal{M}_0$, the identity-connected component of $\mathcal{M}$. In the non-Grassmannian cases $\mathcal{M} = \mathcal{M}_0$, but in the Grassmannian cases $\mathcal{M}$ is a union of disconnected components, each of which is the Cartan immersion of a different G/H' [40].

In order to make connections with subsection 3.5 we consider translations of the Cartan-immersed G/H . In the Grassmannian cases we translate by left-multiplying by the diagonal matrix X . (In the unitary and orthogonal cases if $\det X = -1$ then the resulting construction is no longer a subset of $SU(N)$ or $SO(N)$, but lies in the determinant -1 part of $U(N)$ or $O(N)$, respectively.) In the case of $SU(N)/SO(N)$ we do not need to translate the Cartan construction. For $SU(N)/Sp(N)$ we translate by J (which has determinant 1). In the remaining cases of $SO(2n)/U(n)$ and $Sp(2n)/U(n)$ we translate by iJ , which has determinant $(-1)^n$ and so will be a translation into the determinant -1 part of $O(2n)$ if and only if n is odd. The full set of translated Cartan immersions is then

$$\begin{aligned} \frac{SU(N)}{S(U(M) \times U(N-M))} &\cong \{UXU^\dagger | U \in SU(N)\} \\ \frac{SO(N)}{S(O(M) \times O(N-M))} &\cong \{UXU^t | U \in SO(N)\} \\ \frac{Sp(N)}{Sp(M) \times Sp(N-M)} &\cong \{UXU^{-1} | U \in Sp(N)\} \\ \frac{SU(N)}{SO(N)} &\cong \{U^*U^\dagger | U \in SU(N)\} \\ \frac{SU(N)}{Sp(N)} &\cong \{U^*JU^\dagger | U \in SU(N)\} \\ \frac{SO(2n)}{U(n)} &\cong \{iUJU^t | U \in SO(2n)\} \\ \frac{Sp(2n)}{U(n)} &\cong \{iUJU^{-1} | U \in Sp(2n)\}, \end{aligned}$$

and we treat each of these in turn in appendices 5.2.1 - 5.2.7.

5.2 The Boundary S -matrix Constraints

The aim of this appendix is to show in every case that the above constructions of the symmetric spaces can be described in terms of constraints (those from subsection 3.5) on a single complex $N \times N$ matrix $E \in Gl(N, \mathbb{C})$.

$$5.2.1 \quad \{UXU^\dagger | U \in SU(N)\} = \{E | E^\dagger = E, E^2 = I, Tr(E) = 2M - N\}$$

$$\{UXU^\dagger | U \in SU(N)\} \subseteq \{E | E^\dagger = E, E^2 = I, Tr(E) = 2M - N\} \quad \text{is obvious.}$$

Now if $E^\dagger = E$ then $\exists U \in SU(N)$ s.t. $E = UDU^\dagger$, where D is diagonal.

$$E^2 = I \Rightarrow D^2 = I \text{ so } D \text{ has diagonal entries } \pm 1$$

Thus, after possible reordering of the diagonal entries (which we absorb into U)

$$D = \begin{pmatrix} I_{\tilde{M}} & 0 \\ 0 & -I_{N-\tilde{M}} \end{pmatrix}$$

The constraint on $Tr(E)$ implies $D = X$ and so we have $E = UXU^\dagger$, as required.

$$5.2.2 \quad \{UXU^t | U \in SO(N)\} = \{E | E^\dagger = E, E^2 = I, E^t = E, Tr(E) = 2M - N\}$$

$$\{UXU^t | U \in SO(N)\} \subseteq \{E | E^\dagger = E, E^2 = I, E^t = E, Tr(E) = 2M - N\} \quad \text{is obvious.}$$

Now $E^\dagger = E$ and $E^t = E$ imply that E is a real symmetric matrix, therefore

$$\exists U \in SO(N) \text{ s.t. } E = UDU^t, \text{ where } D \text{ is diagonal}$$

As in the case above the conditions $E^2 = I$ and $Tr(E) = 2M - N$ require that $D = X$, so we have $E = UXU^t$, as required.

$$5.2.3 \quad \{UXU^{-1} | U \in Sp(N)\} = \{E | E^\dagger = E, E^2 = I, JE^t J = -E, Tr(E) = 2M - N\}$$

$$\{UXU^{-1} | U \in Sp(N)\} \subseteq \{E | E^\dagger = E, E^2 = I, JE^t J = -E, Tr(E) = 2M - N\} \quad \text{is obvious.}$$

In order to show the converse we consider the following correspondence¹³. Consider the representation of $\mathbb{H}$ by the 2×2 complex matrices

$$\mathbf{a} = a_0 \mathbf{e} + a_1 \mathbf{i} + a_2 \mathbf{j} + a_3 \mathbf{k} \mapsto \mathbb{A} = a_0 \mathbb{E} + a_1 \mathbb{I} + a_2 \mathbb{J} + a_3 \mathbb{K}$$

¹³Thanks to Ian McIntosh for providing this suggestion.

where

$$\mathbb{E} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \mathbb{I} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \quad \mathbb{J} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \mathbb{K} = \begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix}$$

If we define complex conjugation on $\mathbb{H}$ in the standard way (note that this corresponds to hermitian conjugation of the matrices), and consider quaternions of unit length

$$\mathbf{a}\mathbf{a}^* = (a_0^2 + a_1^2 + a_2^2 + a_3^2)\mathbf{e} = \mathbf{e}$$

these correspond to elements of $SU(2) = Sp(2)$ under the map. (Recall our definition of $Sp(2n)$, as being the subset of $U(2n)$ satisfying the condition $AJA^t = J$. In the $n = 1$ case this is simply the constraint $\det \mathbb{A} = 1$, and so $SU(2) = Sp(2)$.)

This correspondence can be generalized to $Sp(2n)$ in the following way. Consider

$$\mathcal{A} = \begin{pmatrix} \mathbf{a}_{11} & \mathbf{a}_{12} & \dots & \mathbf{a}_{1n} \\ \mathbf{a}_{21} & \mathbf{a}_{22} & \dots & \mathbf{a}_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{a}_{n1} & \mathbf{a}_{n2} & \dots & \mathbf{a}_{nn} \end{pmatrix} \mapsto \left(\begin{array}{c|c|c|c} \mathbb{A}_{11} & \mathbb{A}_{12} & \dots & \mathbb{A}_{1n} \\ \hline \mathbb{A}_{21} & \mathbb{A}_{22} & \dots & \mathbb{A}_{2n} \\ \hline \vdots & \vdots & \ddots & \vdots \\ \hline \mathbb{A}_{n1} & \mathbb{A}_{n2} & \dots & \mathbb{A}_{nn} \end{array} \right) = A$$

where the quaternion, $\mathbf{a}_{ij} \mapsto \mathbb{A}_{ij}$, the 2×2 block, in the way described above. We define

$$J = \left(\begin{array}{c|c|c|c} \mathbb{J} & 0 & \dots & 0 \\ \hline 0 & \mathbb{J} & & 0 \\ \hline \vdots & & \ddots & \\ \hline 0 & 0 & & \mathbb{J} \end{array} \right) \quad \text{where } \mathbb{J} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix};$$

then the conditions $\mathcal{A}\mathcal{A}^\dagger = \mathcal{I}$ (where $\mathcal{I}$ is the quaternionic identity matrix) and $A \in Sp(2n)$ are in exact correspondence.

We now appeal to the fact that the (quaternionic) unitary matrix $\mathcal{A}$ can be diagonalized

$$\mathcal{Q}^\dagger \mathcal{A} \mathcal{Q} = \mathcal{D} \quad \text{where } \mathcal{Q}, \mathcal{D} \in U(n, \mathbb{H}) \text{ and } \mathcal{D} \text{ is diagonal}$$

Under the isomorphism we have established between $U(n, \mathbb{H})$ and $Sp(2n)$ this statement corresponds to $Q^\dagger A Q = D$ where $Q, D \in Sp(2n)$ and D is of the form

$$D = \left(\begin{array}{c|c|c|c} \mathbb{D}_1 & 0 & & \\ \hline 0 & \mathbb{D}_2 & \ddots & \\ \hline & \ddots & \ddots & 0 \\ \hline & & 0 & \mathbb{D}_n \end{array} \right) \quad \text{with } \mathbb{D}_i \in SU(2).$$

We can diagonalize each $SU(2)$ block $\mathbb{D}_i$ by conjugating by some $\mathbb{P}_i \in SU(2)$. If we form the matrix P by placing these $SU(2)$ blocks, in order, down the diagonal then we have

$$P^\dagger D P = X \quad \text{where } X \text{ is diagonal.}$$

Since $SU(2) = Sp(2)$ we find that $P \in Sp(2n)$. Thus, for $A \in Sp(2n)$, we have obtained the conjugation

$$Q^\dagger P^\dagger A P Q = X \quad \text{where } Q, P, X \in Sp(2n) \text{ and } X \text{ is diagonal.}$$

Taking $U = PQ$, we have shown the following to be true:

For all $A \in Sp(2n) \exists U \in Sp(2n)$ s.t. $U^{-1} A U = X$ where X is diagonal.

Now, recalling the set $\{E | E^\dagger = E, E^2 = I, J E^t J = -E, \text{Tr}(E) = 2M - N\}$, we have

$$E^2 = I \quad \text{and} \quad J E^t J = -E \Rightarrow E J E^t = J.$$

Thus $E \in Sp(N)$ and we apply our result above to give $E = U X U^{-1}$ for some $U \in Sp(N)$. We find that $X^2 = I$ and $\text{Tr}(X) = 2M - N$, so we have

$$X = \begin{pmatrix} I_M & 0 \\ 0 & -I_{N-M} \end{pmatrix}.$$

(Note: any X satisfying the above is conjugate to this X via some $Sp(2n)$ matrix, which we absorb into U .) Thus we have $E = U X U^{-1}$ with the required X .

$$5.2.4 \quad \{U^* U^\dagger | U \in SU(N)\} = \{E | E^\dagger E = I, E^t = E, \det E = 1\}$$

$$\{U^* U^\dagger | U \in SU(N)\} \subseteq \{E | E^\dagger E = I, E^t = E, \det E = 1\} \quad \text{is obvious.}$$

Now if $E^\dagger E = I$ then $\exists Q \in SU(N)$ s.t. $E = Q D Q^\dagger$, where D is diagonal.

If we impose the condition $E^t = E$, we have

$$Q^* D Q^t = Q D Q^\dagger \Rightarrow D Q^t Q = Q^t Q D.$$

Since D is diagonal, we can find a diagonal matrix C s.t. $C^2 = D$ and with the property $C Q^t Q = Q^t Q C$. Thus, we have

$$\begin{aligned} E &= Q C^2 Q^\dagger \\ &= Q^* Q^t Q C^2 Q^\dagger \\ &= Q^* C Q^t Q C Q^\dagger \\ &= U^* U^\dagger \quad \text{setting } Q^* C Q^t = U^*. \end{aligned}$$

We see that $\det U = \det C^* = \pm 1$, and so we have $U \in \{V \in U(N) \mid \det V = \pm 1\}$.

Next, we show that the two sets $\{U^*U^\dagger \mid UU^\dagger = I, \det U = 1\}$ and $\{U^*U^\dagger \mid UU^\dagger = I, \det U = -1\}$ are in fact the same. This is trivially true for $N = 1$, so we assume $N \geq 2$ and suppose we have an element E belonging to the first set, that is $E = U^*U^\dagger$ for $U \in SU(N)$. We consider $U' = UT$ where

$$T = \left(\begin{array}{cc|c} 0 & 1 & 0 \\ 1 & 0 & \\ \hline 0 & & I_{N-2} \end{array} \right) \quad (\text{note: } T \in O(N), \det T = -1),$$

and we see that

$$\begin{aligned} U'^*U'^\dagger &= U^*TT^tU^\dagger \\ &= U^*U^\dagger \\ &= E. \end{aligned}$$

We note that $U'^*U'^\dagger$ is a member of the second set, right multiplying by T in this way is a self-inverse operation, and thus we have shown the two sets to be equal.

Thus, we have established the equality

$$\{U^*U^\dagger \mid U \in SU(N)\} = \{E \mid E^\dagger E = I, E^t = E, \det E = 1\},$$

as required for section 3.5.4.

$$5.2.5 \quad \{U^*JU^\dagger \mid U \in U(N), \det U = \pm 1\} = \{E \mid E^\dagger E = I, E^t = -E, \det E = 1\}$$

$$\{U^*JU^\dagger \mid U \in U(N), \det U = \pm 1\} \subseteq \{E \mid E^\dagger E = I, E^t = -E, \det E = 1\} \quad \text{is obvious.}$$

Now if $E^\dagger E = I$ then $\exists Q \in SU(N)$ s.t. $E = QDQ^\dagger$, where D is diagonal.

If we impose the condition $E^t = -E$, we have

$$Q^*DQ^t = -QDQ^\dagger \Rightarrow DQ^tQ = -Q^tQD.$$

We denote the diagonal entries of D by D_i , so that D has entries $D_i\delta_{ij}$. If we denote the entries of Q^tQ by P_{ij} then the condition above becomes

$$D_i\delta_{ij}P_{jk} = -P_{ij}D_j\delta_{jk} \Rightarrow D_iP_{ik} = -D_kP_{ik}.$$

We see that $P_{ik} \neq 0 \Rightarrow D_i = -D_k$. By a suitable rearrangement of the diagonal entries of D (which we absorb by redefining Q) we take D to be of the form

$$D = \left(\begin{array}{c|c|c|c|c} d_1 I_{n_1} & 0 & \dots & \dots & 0 \\ \hline 0 & -d_1 I_{m_1} & 0 & & \vdots \\ \hline \vdots & 0 & d_2 I_{n_2} & \ddots & \vdots \\ \hline \vdots & & \ddots & \ddots & 0 \\ \hline 0 & \dots & \dots & 0 & -d_k I_{m_k} \end{array} \right) \quad \text{where } \begin{array}{l} n_i \geq m_i \geq 0, \\ n_i \geq 1, \\ d_j \neq \pm d_i \text{ if } j \neq i. \end{array}$$

Then $Q^t Q$ (which is symmetric) must have the form

$$Q^t Q = \left(\begin{array}{c|c|c|c|c|c|c} 0 & P_1 & 0 & 0 & \dots & 0 & 0 \\ \hline P_1^t & 0 & 0 & 0 & \dots & 0 & 0 \\ \hline 0 & 0 & 0 & P_2 & & \vdots & \vdots \\ \hline 0 & 0 & P_2^t & 0 & & \vdots & \vdots \\ \hline \vdots & \vdots & & & \ddots & \vdots & 0 \\ \hline 0 & 0 & \dots & \dots & \dots & 0 & P_k \\ \hline 0 & 0 & \dots & \dots & 0 & P_k^t & 0 \end{array} \right) \quad \text{where } P_i \text{ is of size } n_i \text{ rows by } m_i \text{ columns.}$$

We recall that $Q^t Q \in SU(N)$, so that the n_i rows of P_i are orthonormal with respect to the inner product on $\mathbb{C}^{m_i}$. Consequently, $m_i \geq n_i$ and so we must have $n_i = m_i$, with $P_i \in SU(n_i)$, for all i .

We now decompose D into two diagonal matrices $D = \tilde{D} \mathcal{E}$ where

$$\tilde{D} = \left(\begin{array}{c|c|c|c} -id_1 I_{2n_1} & 0 & \dots & 0 \\ \hline 0 & -id_2 I_{2n_2} & & \vdots \\ \hline \vdots & & \ddots & 0 \\ \hline 0 & \dots & 0 & -id_k I_{2n_k} \end{array} \right), \quad \mathcal{E} = \left(\begin{array}{c|c|c} iI_{n_1} & 0 & \\ \hline 0 & -iI_{n_1} & \\ \hline & \ddots & \\ \hline & & iI_{n_k} & 0 \\ & & 0 & -iI_{n_k} \end{array} \right).$$

These matrices satisfy

$$\tilde{D} Q^t Q = Q^t Q \tilde{D} \quad \mathcal{E} Q^t Q = -Q^t Q \mathcal{E}.$$

Recall $\det E = 1 \Rightarrow \det D = 1$, so since $\det \mathcal{E} = 1$ we also have $\det \tilde{D} = 1$. We now choose a diagonal matrix C such that $C^2 = \tilde{D}$ and $C Q^t Q = Q^t Q C$ whose determinant will be ± 1 . (Note that all possible C will have the same determinant.) Now we consider

$$\begin{aligned} E &= Q D Q^\dagger \\ &= Q C^2 \mathcal{E} Q^\dagger \\ &= Q^* Q^t Q C \mathcal{E} C Q^\dagger \quad C \mathcal{E} = \mathcal{E} C \text{ as } C \text{ and } \mathcal{E} \text{ are diagonal} \\ &= Q^* C Q^t Q \mathcal{E} C Q^\dagger. \end{aligned}$$

We set $R = Q\mathcal{E}Q^\dagger$, then $Q\mathcal{E} = RQ$ so that $E = Q^*CQ^tRQCQ^\dagger$. Now we consider the properties of R

$$\begin{aligned} R^\dagger &= Q\mathcal{E}^\dagger Q^\dagger = -Q\mathcal{E}Q^\dagger = -R \\ R^t &= Q^*\mathcal{E}Q^t = Q^*\mathcal{E}Q^tQQ^\dagger = -Q^*Q^tQ\mathcal{E}Q^\dagger = -R \Rightarrow R^* = R \\ R^2 &= Q\mathcal{E}Q^\dagger Q\mathcal{E}Q^\dagger = -I \quad \text{as } \mathcal{E}^2 = -I. \end{aligned}$$

Further, $\det R = 1$ so we have $R \in SO(N)$ and $R^2 = -I$. Thus, appealing to the argument contained in the next section 5.2.6, we can find a matrix $O \in O(N)$ such that $R = OJO^t$. Substituting this into our expression for E we have $E = Q^*CQ^tOJO^tQCQ^\dagger$. Setting $U^* = Q^*CQ^tO \rightarrow U^\dagger = O^tQCQ^\dagger$ we have

$$E = U^*JU^\dagger \quad \text{where } U \in U(N), \det U = \pm 1.$$

We have shown $\{U^*JU^\dagger | U \in U(N), \det U = \pm 1\} = \{E | E^\dagger E = I, E^t = -E, \det E = 1\}$, as required. However, unlike the last subsection 5.2.4, here the $\det U = \pm 1$ subsets are different, for consider:

suppose $U^*JU^\dagger = V^*JV^\dagger$ where $U, V \in U(N), \det U = 1, \det V = -1 \Rightarrow \det(V^tU^*) = -1$:
then $U^*JU^\dagger = V^*JV^\dagger \Rightarrow V^tU^*JU^\dagger V = J \Rightarrow V^tU^* \in Sp(N) \Rightarrow \det(V^tU^*) = 1 \quad \times$

The subsets of $U(N)$ such that $\det = \pm 1$ are isomorphic via multiplication by ω where $\omega^N = -1$. Thus

$$\{U^*JU^\dagger | U \in U(N), \det U = -1\} = \omega^2 \{U^*JU^\dagger | U \in SU(N)\}$$

So we have

$$\{E | E^\dagger E = I, E^t = -E, \det E = 1\} = \{1, \omega^2\} \times \{U^*JU^\dagger | U \in SU(N)\},$$

as required by subsection 3.5.4.

$$5.2.6 \quad \{iUJU^t | U \in O(2n)\} = \{E | E^\dagger = E, E^2 = I, E^t = -E, \}$$

$\{iUJU^t | U \in O(2n)\} \subseteq \{E | E^\dagger = E, E^2 = I, E^t = -E, \}$ is obvious.

Now, $E^\dagger = E, E^t = -E \Rightarrow E^* = -E$, so we consider $F = -iE$. Then F is a real matrix satisfying

$$F^t F = I \quad \text{and} \quad F^2 = -I.$$

Since F is orthogonal there exists a matrix $R \in SO(2n)$ such that $R^t F R$ has the form

$$\left(\begin{array}{c|c|c|c} O_1 & 0 & & \\ \hline 0 & O_2 & \ddots & \\ \hline & \ddots & \ddots & 0 \\ \hline & & 0 & O_n \end{array} \right) \quad \text{where } O_i \in O(2).$$

Since $F^2 = -I$ each $O_i^2 = -I_2$, the only solutions for which $O_i \in O(2)$ are

$$O_i = \pm \epsilon = \pm \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

We note that it is possible to conjugate $-\epsilon$ by the matrix

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \in O(2)$$

to obtain ϵ . Thus we can find a matrix $R' \in O(2)^{\otimes n} \subset O(2n)$ such that $R'^t R^t F R R' = J$. If we set $U = R R'$ then we have

$$E = iUJU^t \quad \text{for } U \in O(2n) \quad \text{as required.}$$

We note that $\{iUJU^t | U \in O(2n)\} \neq \{iUJU^t | U \in SO(2n)\}$, since

$$\begin{aligned} iUJU^t &= iVJV^t \quad \text{for } U, V \in O(N) \\ \Rightarrow V^t UJU^t V &= J \\ \Rightarrow V^t U &\in Sp(2n) \\ \Rightarrow \det U &= \det V. \end{aligned}$$

Both sets $\{iUJU^t | U \in SO(2n)\}$ and $\{iUJU^t | U \in O(2n), \det U = -1\}$ are constructions of the symmetric space

$$\frac{SO(2n)}{U(n)}$$

and so $\{E | E^\dagger = E, E^2 = I, E^t = -E\}$ is isomorphic to two copies of the symmetric space, as stated in section 3.5.2.

$$5.2.7 \quad \{iUJU^{-1} | U \in Sp(2n)\} = \{E | E^\dagger = E, E^2 = I, JE^t J = E\}$$

$$\{iUJU^{-1} | U \in Sp(2n)\} \subseteq \{E | E^\dagger = E, E^2 = I, JE^t J = E\} \quad \text{is obvious.}$$

To establish the converse, we consider the conditions on E

$$\begin{aligned} E^2 = I \text{ and } JE^t J = E &\Rightarrow EJE^t = -J \\ E^\dagger = E \text{ and } E^2 = I &\Rightarrow E^\dagger E = I. \end{aligned}$$

If we let $F = -iE$ then the conditions on F are

$$F^\dagger F = I, \quad F^2 = -I \quad \text{and} \quad FJF^t = J.$$

Thus $F \in Sp(2n)$, so from 5.2.3 we can find a $V \in Sp(2n)$ such that $F = VDV^{-1}$, where D is a diagonal matrix. Since $F^2 = -I$ we must also have $D^2 = -I$, so the entries of D must be $\pm i$. As $D \in Sp(2n)$ we cannot choose the signs of these diagonal entries completely arbitrarily and we find we are restricted to

$$D = \left(\begin{array}{c|c|c|c} \pm \mathbb{I} & 0 & & \\ \hline 0 & \pm \mathbb{I} & \ddots & \\ \hline & \ddots & \ddots & 0 \\ \hline & & 0 & \pm \mathbb{I} \end{array} \right), \quad \text{recalling } \mathbb{I} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$$

and with each $\pm$ freely chosen. However, we have $\mathbb{K} \mathbb{I} \mathbb{K}^\dagger = -\mathbb{I}$ and so we can further conjugate in $Sp(2n)$ (which we absorb into V) to ensure that D has all n $\pm$ signs set to $+$.

Now we notice that $J \in Sp(2n)$ and so, as above, there exists some $W \in Sp(2n)$ such that $D = WJW^{-1}$. If we set $U = VW$ then we have obtained $F = UJU^{-1}$ where $U \in Sp(2n)$. Recalling that $E = iF$, we see that we have obtained

$$E = iUJU^{-1} \quad \text{as required.}$$

6 Appendix: discrete ambiguities in boundary parameters

First, recall the $G \times G$ invariance of the Lagrangian under $g \mapsto g_L g g_R^{-1}$. Of course this should really be $\frac{G \times G}{\mathbf{Z}(G)}$ (where $\mathbf{Z}(G)$ is the centre of G , which is finite), since for $z \in \mathbf{Z}(G)$ we have $g = zg z^{-1}$. The physics literature for the bulk principal chiral model generally is not concerned with this. In the same way we do not explore in the text the ambiguities in our boundary parameters, but we wish here at least to state them precisely, and to prove that they are finite in number.

6.1 Chiral BCs

The ambiguity here arises because there may be non-trivial g_1, g_2 such that $g_1 H g_2^{-1} = H$. This requires

$$\alpha(g_1) h \alpha(g_2^{-1}) = g_1 h g_2^{-1} \quad \forall h \in H,$$

and thence

$$g_1^{-1} \alpha(g_1) h = h g_2^{-1} \alpha(g_2) \quad \forall h \in H.$$

So (by setting $h = e$) we see that

$$g_1^{-1} \alpha(g_1) = g_2^{-1} \alpha(g_2) = k \quad \text{where} \quad k \in \frac{G}{H} \cap C(H),$$

where $C(H) < G$ is the centralizer of H . But, for a symmetric space, H is a maximal Lie subgroup (there is no Lie subgroup of greater dimension which contains H), so $\frac{G}{H} \cap C(H)$ is finite, and so the solutions $g_1 \in Hx$, $g_2 \in Hy$ have $x = y$ (since the Cartan immersion is 1-1) and are also finite in number.

6.2 Non-chiral BCs

Here the potential ambiguity is that there may be non-trivial g_1, g_2 such that $g_1 \frac{G}{H} g_2^{-1} = \frac{G}{H}$. In contrast to the chiral case, we can push g_1 through $\frac{G}{H}$:

$$\begin{aligned} g_1 \{ \alpha(g) g^{-1} | g \in G \} g_2^{-1} &= \{ \alpha(\alpha(g_1) g) g^{-1} | g \in G \} g_2^{-1} \\ &= \{ \alpha(\alpha(g_1) g) (\alpha(g_1) g)^{-1} | g \in G \} \alpha(g_1) g_2^{-1} \\ &= \frac{G}{H} \alpha(g_1) g_2^{-1}. \end{aligned}$$

So the boundary is parametrized by G , up to g_0 such that $\frac{G}{H} g_0 = \frac{G}{H}$. This requires $\alpha(k g_0) = g_0^{-1} k^{-1}$ for all $k \in G/H \hookrightarrow G$, so $k^{-1} \alpha(g_0) = g_0^{-1} k^{-1}$. This must hold for $k = e$, so $g_0 \in \mathcal{M}$ of (5.1), and commutes with every element of $G/H \hookrightarrow G$.

Such g_0 form a group, which must be finite: for suppose not, that its algebra is generated by $\mathbf{k}_0 \subset \mathbf{k}$. Then $[\mathbf{k}_0, \mathbf{k}] = 0$. Also $[[\mathbf{h}, \mathbf{k}_0], \mathbf{k}] \subset [[\mathbf{k}, \mathbf{k}_0], \mathbf{h}] + [[\mathbf{h}, \mathbf{k}], \mathbf{k}_0]$, both of which are empty, so $[\mathbf{h}, \mathbf{k}_0] \subset \mathbf{k}_0$. Thus $[\mathbf{k}_0, \mathbf{g}] \subset \mathbf{k}_0$, and $\mathbf{k}_0$ is an ideal, and is therefore trivial.

Note the specialization (mentioned in the text) when $g_1 = g_2$: the boundary is then parametrized by $\alpha(g_1) g_1^{-1}$ and thus by G/H (again quotiented, here by $\mathbf{Z}(G/H)$, those elements of G/H which commute with all of G/H). It is straightforward to propose a compatible PCM boundary S -matrix, though we do not do so here.

7 Appendix: boundary Yang Baxter and crossing-unitarity calculations

In this section we include a representative selection of the BYBE and crossing-unitarity calculations required to obtain the various constraints on the boundary S -matrices that we have considered in this paper. We hope they will be sufficiently illustrative that the interested reader can perform any calculations not presented here for themselves.

First we list the minimal bulk S -matrices derived in [29]

$$\begin{aligned} SU(N) : & \quad \frac{\sigma_u(\theta)}{(1 - \frac{h\theta}{2i\pi})} \left(\text{---} - \frac{h\theta}{2i\pi} \text{X} \right) \\ SO(N) : & \quad \frac{\sigma_o(\theta)}{(1 - \frac{h\theta}{2i\pi})} \left(\text{---} - \frac{h\theta}{2i\pi} \text{X} + \frac{\frac{h\theta}{2i\pi}}{(\frac{h}{2} - \frac{h\theta}{2i\pi})} \right) (\text{---}) \\ Sp(N) : & \quad \frac{\sigma_p(\theta)}{(1 - \frac{h\theta}{2i\pi})} \left(\text{---} - \frac{h\theta}{2i\pi} \text{X} + \frac{\frac{h\theta}{2i\pi}}{(\frac{h}{2} - \frac{h\theta}{2i\pi})} \text{X} \right) \end{aligned}$$

where h is the dual Coxeter number and the scalar prefactors σ are ¹⁴

$$\begin{aligned} \sigma_u(\theta) &= \mp \frac{\Gamma(\frac{\theta}{2i\pi} + \frac{1}{h}) \Gamma(\frac{-\theta}{2i\pi})}{\Gamma(\frac{-\theta}{2i\pi} + \frac{1}{h}) \Gamma(\frac{\theta}{2i\pi})} \\ \sigma_o(\theta) &= - \frac{\Gamma(\frac{\theta}{2i\pi} + \frac{1}{h}) \Gamma(\frac{\theta}{2i\pi} + \frac{1}{2}) \Gamma(\frac{-\theta}{2i\pi}) \Gamma(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{h})}{\Gamma(\frac{-\theta}{2i\pi} + \frac{1}{h}) \Gamma(\frac{-\theta}{2i\pi} + \frac{1}{2}) \Gamma(\frac{\theta}{2i\pi}) \Gamma(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{h})} \\ \sigma_p(\theta) &= - \frac{\Gamma(\frac{\theta}{2i\pi} + \frac{1}{h}) \Gamma(\frac{\theta}{2i\pi} + \frac{1}{2}) \Gamma(\frac{-\theta}{2i\pi}) \Gamma(\frac{-\theta}{2i\pi} + \frac{1}{2} + \frac{1}{h})}{\Gamma(\frac{-\theta}{2i\pi} + \frac{1}{h}) \Gamma(\frac{-\theta}{2i\pi} + \frac{1}{2}) \Gamma(\frac{\theta}{2i\pi}) \Gamma(\frac{\theta}{2i\pi} + \frac{1}{2} + \frac{1}{h})} \end{aligned}$$

where Γ is the gamma function, and

$$h = \begin{cases} N & SU(N) \\ N - 2 & SO(N) \\ N + 2 & Sp(N). \end{cases}$$

Note that

$$\bar{\sigma}_u(\theta) = \sigma_u(i\pi - \theta).$$

¹⁴The $\mp$ sign in the prefactor for $SU(N)$ reflects the fact that this sign is an arbitrary choice. However, from consideration of the boundary bootstrap, details of which are outside the scope of this paper, the indications are that a $-$ sign is required for the model with non-conjugating boundary conditions, whilst a $+$ sign is the preferred choice for representation conjugating BCs.

7.1 BYBE calculations

Recall the boundary Yang Baxter equation

$$S_{ij}^{kl}(\theta - \phi) (I_{jm} \otimes K^{ln}(\theta)) S_{mo}^{np}(\theta + \phi) (I_{oq} \otimes K^{pr}(\phi)) = \\ (I_{ij} \otimes K^{kl}(\phi)) S_{jm}^{ln}(\theta + \phi) (I_{mo} \otimes K^{np}(\theta)) S_{oq}^{pr}(\theta - \phi).$$

For clarity of the calculations we introduce the notation

$$u = \frac{h\theta}{2i\pi}, \quad v = \frac{h\phi}{2i\pi}, \quad u_0 = \frac{h}{4}.$$

In any BYBE calculation the scalar prefactors cancel, and we consider here only the matrix part of the equation.

7.1.1 The $SU(N)$ case with K_1 boundary S -matrix

Substituting into the BYBE with the bulk S -matrix for the $SU(N)$ PCM and K_1 gives

$$\left(\begin{array}{c} \text{---} \\ \text{---} \end{array} - (u-v) \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right) \begin{array}{c} \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} - (u+v) \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right) \begin{array}{c} \text{---} \\ \text{---} \end{array} = \\ \begin{array}{c} \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} - (u+v) \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right) \begin{array}{c} \text{---} \\ \text{---} \end{array} \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} - (u-v) \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right)$$

Expanding out and cancelling where possible, we are left with

$$\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array}$$

Thus, for the equation to be satisfied we require the condition :-

$$\begin{array}{c} \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} = \alpha \begin{array}{c} \text{---} \\ \text{---} \end{array} \quad \text{for some constant } \alpha.$$

7.1.2 The $Sp(N)$ case with K_2 boundary S -matrix

Substituting into the BYBE with the bulk S -matrix for the $Sp(N)$ PCM and K_2 gives

$$\left(\begin{array}{c} \text{---} \\ \text{---} \end{array} - (u-v) \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} + t(u-v) \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right) \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} + \tilde{c}u \begin{array}{c} \text{---} \\ \text{---} \end{array} \right) \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} - (u+v) \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} + t(u+v) \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right) \times \\ \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} + \tilde{c}v \begin{array}{c} \text{---} \\ \text{---} \end{array} \right) = \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} + \tilde{c}v \begin{array}{c} \text{---} \\ \text{---} \end{array} \right) \times \\ \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} - (u+v) \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} + t(u+v) \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right) \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} + \tilde{c}u \begin{array}{c} \text{---} \\ \text{---} \end{array} \right) \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} - (u-v) \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} + t(u-v) \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \right)$$

where $t(u) = \frac{u}{2u_0 - u}$ and $\tilde{c}$ is related to the original S -matrix constant c by the relation $\tilde{c} = \frac{2i\pi c}{h}$. Expanding out, cancelling where possible (noting that the terms involving $\begin{array}{c} \text{---} \\ \text{---} \end{array} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array}$

and  cancel after some simple algebra) and rearranging (some less trivial algebra!) we are left with

$$\begin{aligned}
& -\tilde{c}uv t(u-v)t(u+v)(2 + \tilde{c} \text{ (circle) }) \left(\text{ (cross with circle) } - \text{ (cross with circle) } \right) \\
& + 2\tilde{c}uv t(u-v)t(u+v) \left(\text{ (cross with circle) } - \text{ (cross with circle) } \right) \\
& \tilde{c}^2 uv(u-v) \left(\text{ (cross with circle) } - \text{ (cross with circle) } \right) + \tilde{c}^2 uv t(u-v) \left(\text{ (cross with circle) } - \text{ (cross with circle) } \right) \\
& - \tilde{c}^2 uv(u+v)t(u-v) \left(\text{ (cross with circle) } - \text{ (cross with circle) } \right) \\
& - \tilde{c}^2 uv(u-v)t(u+v) \left(\text{ (cross with circle) } - \text{ (cross with circle) } \right) = 0.
\end{aligned}$$

In order for this to hold we are forced to have

$$\begin{aligned}
\text{ (circle) } &= \alpha \text{ (line) } \quad \text{for some constant } \alpha \\
\text{ (cross with circle) } &= \beta \text{ (cross with circle) } \quad \text{i.e.} \quad (\text{ (circle) })^t = \beta \text{ (cross with circle) }
\end{aligned}$$

(Note: then $\beta^2 = 1$.) We find that the equation is then satisfied provided

$$2\beta - 2 - \tilde{c} \text{ (circle) } = 0.$$

Since $\beta = \pm 1$ we must have :-

$$\begin{aligned}
\text{ (circle) } &= \alpha \text{ (line) } \quad \text{and} \\
(\text{ (circle) })^t &= \text{ (cross with circle) } \iff \tilde{c} \text{ (circle) } = 0 \quad \text{or} \quad (\text{ (circle) })^t = -(\text{ (cross with circle) }) \iff \tilde{c} \text{ (circle) } = -4.
\end{aligned}$$

7.1.3 The $SU(N)$ -conjugating case

Recall the conjugated BYBE

$$\begin{aligned}
S_{ij}^{kl}(\theta - \phi)(I_{jm} \otimes K^{l\bar{n}}(\theta))S_{m\bar{o}}^{\bar{n}p}(\theta + \phi)(I_{\bar{o}q} \otimes K^{p\bar{r}}(\phi)) = \\
(I_{ij} \otimes K^{k\bar{l}}(\phi))S_{j\bar{m}}^{\bar{l}n}(\theta + \phi)(I_{\bar{m}\bar{o}} \otimes K^{n\bar{p}}(\theta))S_{\bar{o}q}^{\bar{p}\bar{r}}(\theta - \phi).
\end{aligned}$$

The K_2 boundary S -matrix does not satisfy the above equation, but K_1 does, under some constraints. Substituting in and using our simplifying notation we get

$$\begin{aligned}
& \left(\text{ (line) } - (u-v) \text{ (cross) } \right) \text{ (circle) } \left(\text{ (line) } - (2u_0 - u - v) \text{ (cross) } \right) \text{ (circle) } = \\
& \text{ (circle) } \left(\text{ (line) } - (2u_0 - u - v) \text{ (cross) } \right) \text{ (circle) } \left(\text{ (line) } - (u-v) \text{ (cross) } \right)
\end{aligned}$$

Expanding out and cancelling all possible terms leaves

$$\text{ (cross with circle) } \text{ (circle) } = \text{ (circle) } \text{ (cross with circle) }$$

So to satisfy the conjugated BYBE we must have $(-\circ-)^t = \pm -\circ-$.

There are two other BYBEs to consider in addition to the $V \otimes V \rightarrow \bar{V} \otimes \bar{V}$ case considered so far. The $\bar{V} \otimes \bar{V} \rightarrow V \otimes V$ BYBE will be similar to the above, so that if we denote by $-\bullet-$ the matrix part of the $\bar{V} \rightarrow V$ boundary S -matrix then we must have $(-\bullet-)^t = \pm -\bullet-$. The last case to consider is $V \otimes \bar{V} \rightarrow \bar{V} \otimes V$, where the BYBE is

$$S_{ij}^{k\bar{l}}(\theta - \phi) \left(I_{jm} \otimes K^{\bar{l}n}(\theta) \right) S_{mo}^{np}(\theta + \phi) (I_{oq} \otimes K^{p\bar{r}}(\phi)) = \\ \left(I_{\bar{i}\bar{j}} \otimes K^{k\bar{l}}(\phi) \right) S_{\bar{j}\bar{m}}^{\bar{l}\bar{n}}(\theta + \phi) (I_{\bar{m}\bar{o}} \otimes K^{\bar{n}p}(\theta)) S_{oq}^{p\bar{r}}(\theta - \phi).$$

Substituting into this, again with simplified notation, we get

$$\left(\begin{array}{c} \text{---} \\ \text{---} \end{array} \right) \left(- (2u_0 - u + v) \times \right) \left(\begin{array}{c} \bullet \\ \text{---} \end{array} \right) \left(\text{---} - (u + v) \times \right) \left(\begin{array}{c} \circ \\ \text{---} \end{array} \right) = \\ \left(\begin{array}{c} \circ \\ \text{---} \end{array} \right) \left(\text{---} - (u + v) \times \right) \left(\begin{array}{c} \bullet \\ \text{---} \end{array} \right) \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} \right) \left(- (2u_0 - u + v) \times \right).$$

Expanding out and cancelling all the terms we can we have

$$\left(\begin{array}{c} \bullet \\ \text{---} \end{array} \right) \left(\begin{array}{c} \circ \\ \text{---} \end{array} \right) - (2u_0 - u + v) \left(\begin{array}{c} \bullet \\ \text{---} \end{array} \right) \left(\begin{array}{c} \circ \\ \text{---} \end{array} \right) - (u + v) \left(\begin{array}{c} \bullet \\ \text{---} \end{array} \right) \left(\begin{array}{c} \circ \\ \text{---} \end{array} \right) = \\ \left(\begin{array}{c} \circ \\ \text{---} \end{array} \right) \left(\begin{array}{c} \bullet \\ \text{---} \end{array} \right) - (2u_0 - u + v) \left(\begin{array}{c} \circ \\ \text{---} \end{array} \right) \left(\begin{array}{c} \bullet \\ \text{---} \end{array} \right) - (u + v) \left(\begin{array}{c} \circ \\ \text{---} \end{array} \right) \left(\begin{array}{c} \bullet \\ \text{---} \end{array} \right).$$

This equation is satisfied provided

$$-\circ-\bullet = -\bullet-\circ = \alpha \text{---} \quad \text{and} \quad \begin{array}{c} \circ \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \circ \end{array} = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \bullet \end{array} = \beta \text{---}.$$

Since $(-\circ-)^t = \pm -\circ-$ and $(-\bullet-)^t = \pm -\bullet-$, we have $\beta = \pm \alpha$ and so have only one independent parameter. The conditions imposed for the conjugated $SU(N)$ case are thus

$$(-\circ-)^t = \pm -\circ- \quad \text{and} \quad (-\bullet-)^t = \pm -\bullet- \quad \text{and} \quad -\circ-\bullet = -\bullet-\circ = \alpha \text{---}.$$

7.2 Crossing-unitarity calculations

Recall the crossing-unitarity equation

$$K^{ij}(\frac{i\pi}{2} - \theta) = S_{jk}^{i\bar{l}}(2\theta) K^{\bar{l}k}(\frac{i\pi}{2} + \theta).$$

We note that it is $S_{jk}^{i\bar{l}}(2\theta)$ that is required here, which is the crossed S -matrix. We obtain it by taking the standard S -matrix substituting $i\pi - 2\theta$ for 2θ and turning the matrix diagrams through 90° . (We note that the process of crossing doesn't alter the S -matrix for the $SO(N)$ and $Sp(N)$ cases, but we go through the process anyway to illustrate the $SU(N)$ cases.) We again make use of some simplifying notation.

7.2.1 The $SO(N)$ case with K_2 boundary S -matrix

Substituting into the crossing-unitarity equation we have

$$\begin{aligned} \frac{\tau(\frac{i\pi}{2} - \theta)}{(1 - c(\frac{i\pi}{2} - \theta))} \left(\right) + \tilde{c}(u_0 - u) \rangle \rangle &= \frac{\sigma_o(i\pi - 2\theta) \tau(\frac{i\pi}{2} + \theta)}{(1 - 2u_o + 2u)(1 - c(\frac{i\pi}{2} + \theta))} \times \\ &\quad \left(\right) (-2(u_0 - u) \times + \frac{u_0 - u}{u} \text{---}) \left(\right) + \tilde{c}(u_0 + u) \rangle \rangle \\ \Rightarrow \frac{\tau(\frac{i\pi}{2} - \theta)}{\tau(\frac{i\pi}{2} + \theta)} \left(\right) + \tilde{c}(u_0 - u) \rangle \rangle &= \frac{\sigma_o(i\pi - 2\theta) \times}{(1 - c(\frac{i\pi}{2} - \theta))} \frac{(1 - 2u_o + 2u) \times}{(1 - c(\frac{i\pi}{2} + \theta))} \left(\begin{array}{c} \left(\begin{array}{c} N + \tilde{c} \bigcirc (u_0 + u) \\ -2(u_0 - u) + \frac{u_0 - u}{u} \end{array} \right) \rangle \\ -2\tilde{c}(u_0 - u)(u_0 + u) \times \diamond \\ + \frac{\tilde{c}(u_0 - u)(u_0 + u)}{u} \rangle \end{array} \right) \end{aligned}$$

In order for this to be satisfied it is necessary to impose $(-\ominus)^t = \pm \ominus$. Considering coefficients of the $\rangle$ terms we find

$$\frac{\tau(\frac{i\pi}{2} - \theta)}{\tau(\frac{i\pi}{2} + \theta)} = \frac{\sigma_o(i\pi - 2\theta)(u_0 + u)}{(1 - 2u_o + 2u)} \left(\frac{1}{u} \mp 2 \right) \frac{(1 - c(\frac{i\pi}{2} - \theta))}{(1 - c(\frac{i\pi}{2} + \theta))}$$

For the coefficients of the $\rangle$ terms to be consistent with this we require a constraint on $\bigcirc$ which depends on the choice of $\pm$, altogether we have the matrix constraints

$$(-\ominus)^t = \ominus \iff \tilde{c} \bigcirc = -4 \quad \text{or} \quad (-\ominus)^t = -(\ominus) \iff \tilde{c} \bigcirc = 0.$$

From the crossing symmetry of the S -matrix we can simplify the constraint on the scalar prefactor, obtaining

$$\frac{\tau(\frac{i\pi}{2} - \theta)}{\tau(\frac{i\pi}{2} + \theta)} = \frac{(u_0 + u)(1 \mp 2u)(1 - c(\frac{i\pi}{2} - \theta))}{(u_0 - u)(1 - 2u)(1 - c(\frac{i\pi}{2} + \theta))} \sigma_o(2\theta)$$

which can be written as

$$\frac{\tau(\frac{i\pi}{2} - \theta)}{\tau(\frac{i\pi}{2} + \theta)} = \begin{cases} \left[\frac{h}{2} \right] \left[\frac{h}{ci\pi} - \frac{h}{2} \right] \sigma_o(2\theta) & (-\ominus)^t = \ominus \\ -[1] \left[\frac{h}{2} \right] \left[\frac{h}{ci\pi} - \frac{h}{2} \right] \sigma_o(2\theta) & (-\ominus)^t = -\ominus. \end{cases}$$

7.2.2 The $SU(N)$ -conjugating case

The K -matrix for $SU(N)$ -conjugating must satisfy a conjugated version of the crossing-unitarity equation,

$$K^{i\bar{j}}(\frac{i\pi}{2} - \theta) = S_{jk}^{il}(2\theta) K^{l\bar{k}}(\frac{i\pi}{2} + \theta).$$

For this case it is not the crossed S -matrix we require, but the standard S -matrix. Substituting in, we have

$$\rho(\frac{i\pi}{2} - \theta) \mathfrak{J} = \frac{\sigma_u(2\theta)}{(1-2u)} \left(\text{---} - 2u \text{X} \right) \rho(\frac{i\pi}{2} + \theta) \mathfrak{J}.$$

On expanding this, we see that $(-\circ-)^t = \pm \text{---}\circ$ is required. Then the condition on the scalar prefactor becomes

$$\frac{\rho(\frac{i\pi}{2} - \theta)}{\rho(\frac{i\pi}{2} + \theta)} = \frac{(1 \mp 2u)}{(1-2u)} \sigma_u(2\theta),$$

which can be written as

$$\frac{\rho(\frac{i\pi}{2} - \theta)}{\rho(\frac{i\pi}{2} + \theta)} = \begin{cases} \sigma_u(2\theta) & (-\circ-)^t = \text{---}\circ \\ -[1]\sigma_u(2\theta) & (-\circ-)^t = -\text{---}\circ. \end{cases}$$

8 Appendix: unitarity and hermitian analyticity calculations

In this section we prove the two results, concerning complex parameters that could consistently be set to 1, stated in section 3.2.

8.1 The non-conjugating case

We start from (3.8),

$$E^\dagger = E \quad \text{and} \quad E^2 = \alpha I \quad \text{where} \quad \alpha \in U(1).$$

Since E is hermitian we can diagonalize it as

$$D = QEQ^\dagger \quad \text{where} \quad Q \in SU(N).$$

Then we have

$$D^\dagger = QE^\dagger Q^\dagger = QEQ^\dagger = D \Rightarrow D^* = D.$$

Further,

$$D^2 = QEQ^\dagger QEQ^\dagger = QE^2 Q^\dagger = \alpha I.$$

Now $D^* = D \Rightarrow \alpha \in \mathbb{R}^+$ so $\alpha = 1$, as stated in 3.2.1.

8.2 The conjugating case

We start from (3.13)

$$\begin{aligned} \alpha E &= F^\dagger & \text{and} & & \rho(\theta) &= \alpha \omega(-\theta^*)^* \\ EF &= \beta I & \text{and} & & \rho(\theta) \omega(-\theta) &= \frac{1}{\beta}. \end{aligned}$$

We can use the rescaling freedom in $K(\theta) = \rho(\theta)E$ and $K'(\theta) = \omega(\theta)F$,

$$\begin{aligned} E &\rightarrow \lambda E, & \rho(\theta) &\rightarrow \frac{1}{\lambda} \rho(\theta), \\ F &\rightarrow \kappa F, & \omega(\theta) &\rightarrow \frac{1}{\kappa} \omega(\theta), \end{aligned}$$

to set both α and β to 1. Once this has been done some rescaling freedom still remains: the phase shift $\lambda = e^{i\psi}$, $\kappa = e^{-i\psi}$ leaves the constraints unchanged. Thus, we can also insist that $\det E = 1 \iff \det F = 1$ as stated in 3.2.2.

References

- [1] A. Belavin and V. Drinfeld, *Solutions of the classical Yang-Baxter equation for simple Lie algebras*, Funct. Anal. **17** (1983) 220
- [2] V. Drinfeld, *Hopf algebras and the quantum Yang-Baxter equation*, Sov. Math. Dokl. **32** (1985) 254
- [3] J. M. Evans, M. Hassan, N. J. MacKay and A. Mountain, *Conserved charges and supersymmetry in principal chiral models*, hep-th/9711140
- [4] J. M. Evans, M. Hassan, N. J. MacKay and A. Mountain, *Local conserved charges in principal chiral models*, Nucl. Phys. **B561** (1999) 385, hep-th/9902008
- [5] J. M. Evans, *Integrable sigma-models and Drinfeld-Sokolov hierarchies*, hep-th/0101231, *From PCM to KdV and back*, hep-th/0103250
- [6] N. J. MacKay, *The $SO(N)$ principal chiral field on a half-line*, J. Phys. **A32** (1999) L189, hep-th/9903137
- [7] I. V. Cherednik, *Factorizing particles on a half-line and root systems*, Theor. Math. Phys. **61** (1983) 977
- [8] N. J. MacKay, *Fusion of $SO(N)$ reflection matrices*, Phys. Lett. **B357** (1995) 89, hep-th/9505124
- [9] J. Abad and M. Rios, *Non-diagonal solutions to reflection equations in $su(n)$ spin chains*, Phys. Lett. **B352** (1995), 92, hep-th/9502129
- [10] E. Corrigan, *Reflections*, hep-th/9601055
- [11] G. M. Gandenberger, *On $a_2^{(1)}$ reflection matrices and affine Toda theories*, Nucl. Phys. **B542**(1999), 659, hep-th/9806003, and
New non-diagonal solutions to the $a_n^{(1)}$ boundary Yang-Baxter equation, hep-th/9911178
- [12] H. J. de Vega and A. González-Ruiz, *Boundary K -matrices for the six-vertex and the $n(2n-1)$ A_{n-1} vertex models*, J. Phys. **A26** (1993), L519, hep-th/9211114, and
Boundary K -matrices for the XYZ, XXZ and XXX spin chains, J. Phys. **A27** (1994), 6129, hep-th/9306089
- [13] A. Lima-Santos, *Reflection K -matrices for 19-vertex models*, Nucl. Phys. **B558** (2000), 637, solv-int/9906003
- [14] M. Lüscher, *Quantum non-local charges and the absence of particle production in the 2D non-linear σ -model*, Nucl. Phys. **B135** (1978) 1
- [15] D. Bernard, *Hidden Yangians in 2D massive current algebras*, Commun. Math. Phys. **137** (1991) 191
- [16] P. E. Dorey, *Root systems and purely elastic S -matrices*, Nucl. Phys. **B358** (1991) 654.

- [17] S. Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Academic Press 1978 (2nd. ed.)
- [18] A. T. Fomenko, *The Plateau problem*, vols. I and II, Gordon and Breach 1990
- [19] M. F. Mourad and R. Sasaki, *Non-linear sigma models on a half plane*, Int. J. Mod. Phys. **A11** (1996) 3127, [hep-th/9509153](#)
- [20] J. M. Evans and A. J. Mountain, *Commuting charges and symmetric spaces*, Phys. Lett. **B483** (2000) 290, [hep-th/0003264](#)
- [21] M. Kato and T. Okada, *D-branes on group manifolds*, Nucl. Phys. **B499** (1997) 583, [hep-th/9612148](#)
- [22] G. Felder, J. Fröhlich, J. Fuchs and C. Schweigert, *The geometry of WZW branes*, J. Geom. Phys. **34** (2000) 132, [hep-th/9909030](#)
- [23] S. Stanciu, *D-branes on group manifolds*, [hep-th/9909163](#)
- [24] M. Moriconi and A. de Martino, *Quantum integrability of certain boundary conditions*, Phys. Lett. **B447** (1999) 292, [hep-th/9809178](#)
- [25] M. Moriconi and A. de Martino, *Boundary S-matrix for the Gross-Neveu model*, Phys. Lett. **B451** (1999) 354, [hep-th/9812009](#)
- [26] Y. Y. Goldschmidt and E. Witten, *Conservation laws in some two-dimensional models*, Phys. Lett. **B91** (1980), 392
- [27] S. Ghoshal and A. Zamolodchikov, *Boundary S-matrix and boundary state in two-dimensional integrable quantum field theory*, Int. J. Mod. Phys. **A9** (1994), 3841, [hep-th/9306002](#)
- [28] J. Luis Miramontes, *Hermitian analyticity versus real analyticity in two-dimensional factorized S-matrix theories*, [hep-th/9901145](#)
- [29] E. Ogievetsky, N. Reshetikhin, P. Wiegmann, *The principal chiral field in two dimensions on classical lie algebras: the Bethe-ansatz solution and factorized theory of scattering*, Nucl. Phys. **B280** (1987) 45-96
- [30] A. Fring and R. Köberle, *Affine Toda field theory in the presence of reflecting boundaries*, Nucl. Phys. **B419** (1994), 647, [hep-th/9309142](#)
- [31] E. Corrigan and Z.-M. Sheng, *Classical integrability of the $O(N)$ nonlinear sigma model on a half-line*, Int. J. Mod. Phys. **A12** (1997) 2825, [hep-th/9612150](#),
J. Prata, *Reflection factors for the principal chiral model*, Phys. Lett. **B438** (1998) 115, [hep-th/9806222](#)
- [32] C. Klimcik and P. Severa, *Open strings and D-branes in WZNW models*, Nucl. Phys. **B488** (1997) 653, [hep-th/9609112](#),
A. Recknagel and V. Schomerus, *D-branes in Gepner models*, Nucl. Phys. **B531** (1998) 185, [hep-th/9712186](#),
A. Alekseev and V. Schomerus, *D-branes in the WZW model*, Phys. Rev. **D60** (1999) 61901, [hep-th/9812193](#)

- [33] A. Doikou and R. Nepomechie, *Bulk and boundary S-matrices for the $SU(N)$ chain*, Nucl. Phys. **B521** (1998), 547, [hep-th/9803118](#)
- [34] D. Arnaudon, J. Avan, N. Crampé, A. Doikou, L. Frappat, E. Ragoucy, *Classification of reflection matrices related to (super) Yangians and application to open spin chain models*, [math.QA/0304150 v2](#)
- [35] S. Ghoshal, *Boundary S-matrix of the $SO(N)$ -symmetric non-linear sigma model*, Phys. Lett. **B334** (1994), 363, [hep-th/9401008](#)
- [36] M. Moriconi, *Integrable boundary conditions and reflection matrices for the $O(N)$ nonlinear sigma model*, Nucl. Phys. **B619** (2001), 396, [hep-th/0108039](#) and *Integrable boundary conditions for the $O(N)$ nonlinear sigma model*, [hep-th/0111195](#)
- [37] J. Donin and S. Shnider, *Quantum symmetric spaces*, J. Pure. Appl. Algebra **100** (1995), 103, [hep-th/9412031](#),
M. Dijkhuizen, *Some remarks on the construction of quantum symmetric spaces*, [math.QA/9512225](#), and
M. Noumi and T. Sugitani, *Quantum symmetric spaces and related q -orthogonal polynomials*, [math.QA/9503225](#)
- [38] A. Molev, M. Nazarov and G. Ol'shanskii, *Yangians and classical Lie algebras*, Russ. Math. Surveys **51** (1996) 205,
A. Molev and G. Olshanski, *Centralizer construction for twisted Yangians*, [q-alg/9712050](#)
- [39] H. Eichenherr, *More about nonlinear sigma models on symmetric spaces*, Nucl. Phys. **B164** (1980), 528, err.ibid. **B282** (1987), 745
- [40] J.-P. Antoine and B. Piette, *Classical nonlinear sigma models on Grassmann manifolds of compact or noncompact type*, J. Math. Phys. **28** (1987), 2753
- [41] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, OUP 3rd edition – see ch.14 and references therein
P. Fendley, *Integrable sigma models and perturbed coset models*, [hep-th/0101034](#)